

MINUARTIA SABALANICA (CARYOPHYLLACEAE), A NEW SPECIES FROM NW IRAN

G. Mostafavi, M. Assadi, T. Nejadsattari, F. Sharifnia & I. Mehregan

Received 26.02.2011. Accepted for publication 10.08.2011.

Mostafavi, G., Assadi, M. Nejadsattari, T., Sharifnia, F. & Mehregan. I: 2011 12 31: *Minuartia sabalanica* (*Caryophyllaceae*), a new species from NW Iran. –*Iran. J. Bot.* 17 (2): 220-226. Tehran.

A progressive revision of the Iranian material belonging to *Minuartia* L. (*Caryophyllaceae*) led to the recognition of a new species *Minuartia sabalanica* based on the material from NW Iran. The new species is morphologically related to *M. rimarum*, an endemic only known from the central and southern Turkey, and *M. umbellulifera* known from Turkey and Iran. Diagnostic characters of the new species are discussed. In addition, a Latin diagnosis, description and illustration of the new species are given.

Golaleh Mostafavi (correspondence <g.mostafavi@iausr.ac.ir>), Taher Nejadsattari and Iraj Mehregan, Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran. –Mostafa Assadi, Research Institute of Forests and Rangelands, P. O> Box 13185-116, Tehran ,Iran. –Fariba Sharifnia, Department of Biology, North Tehran Branch, Islamic Azad University, Tehran, Iran.

Key words. Taxonomy, *Minuartia*, *Caryophyllaceae*, new species, Iran.

گونه جدید (Caryophyllaceae) *Minuartia sabalanica* از شمال غربی ایران

گلalte مصطفوی، دانشجوی دکتری سیستماتیک گیاهی، دانشگاه آزاد اسلامی، واحد علوم و تحقیقات، گروه زیست شناسی، تهران، ایران.

مصطفی اسدی، استاد پژوهش، موسسه تحقیقات جنگلها و مراتع کشور، تهران، ایران.

طاهر نژادستاری، دانشیار گروه زیست شناسی، دانشگاه آزاد اسلامی، واحد علوم و تحقیقات، تهران، ایران.

فریبا شریف نیا، دانشیار گروه زیست شناسی، دانشگاه آزاد اسلامی، واحد تهران شمال، تهران، ایران.

ایرج مهرگان، استادیار گروه زیست شناسی، دانشگاه آزاد اسلامی، واحد علوم و تحقیقات، تهران، ایران.

مطالعه برای بازنگری جنس *Minuartia* در ایران، منجر به معرفی گونه جدید *Minuartia sabalanica* به دنیای علم شد. این نمونه که از شمال غربی ایران جمع آوری شده، از نظر ریخت شناسی به *M. rimarum* گونه‌ای انحصاری مناطق مرکزی ترکیه، و نیز *M. umbellulifera* که قبلا از ترکیه و ایران شناخته شده بوده، نزدیک است. صفات تشخیصی گونه جدید بحث شده است. هم چنین دیاگنوza لاتین، شرح و تصویر گونه جدید آورده شده است.

INTRODUCTION

The genus *Minuartia* L. (*Caryophyllaceae* - *Alsinoideae*) comprises ca. 120 usually dwarf, annual or perennial herbaceous species throughout the northern hemisphere (Bittrich 1993). Its members, especially the perennial ones grow usually in inhospitable extreme conditions such as rocky ledges, stony soils and alpine environments (McNeill 1962, 1963). It occurs in Europe with over 50, former U.S.S.R with 44 and Turkey with 42 species (Kamari

1997; Halliday 1964; Meikle 1977; Schischkin 1936; McNeill 1967). Parsa (1951) has introduced 24 species for Iran some of which were reduced as synonyms in Flora Iranica (Rechinger 1988). According to Rechinger (1988), Iran with 21 species belonging to six sections, is one of the biodiversity centers of the genus. Although, Assadi (1984) recorded *M. umbellulifera* (Boiss.) McNeill from NW Iran, it was not mentioned in Flora Iranica (Rechinger 1988). More than half of *Minuartia* species in Iran are reported from northwestern part of the country (Rechinger 1988).

Table 1. Comparison of some morphological characters in *Minuartia sabalanica*, *M. umbellulifera* and *M. rimarum*.

| Characters Species | Stem indumentum | Height (cm) | Leaf shape | Leaf indumentum | Leaf veins | Stem leaves length (mm) | Leaf desity flowers | Number of flowers | Petal shape | Pedicel length (mm) | Seed number per capsule |
|---|---|----------------|---|--|---------------|----------------------------------|---------------------------|-------------------------|-------------------------|---------------------------|----------------------------------|
| <i>M. umbellulifera</i> | not hairy | 6.5-11 | setaceous- terete | not hairy | 1 | 3.2-6 | dense | 1-10 | oblateolate | 6-11 | 9-16 |
| <i>M. rimarum</i> var. <i>multiflora</i> | glandular-hairy | up to 8 | linear to linear oblong, straight, flat | densely glandular-hirsute | 3 | ca. 4 | dense | 2-5 | obovate | ca. 5-11 | unknown |
| <i>M. rimarum</i> var. <i>rimarum</i> | glandular-hairy | up to 8 | linear to linear oblong, flat, slightly curved | sub glabrous to moderately glandular | 3 | ca.3.5- 7.5 | almost loose | 1-2 | oblateolate | ca. 5- 14.5 | unknown |
| <i>M. sabalanica</i> | glandular hair on the inflorescence | 7.5-11 | Linear, canaliculate, flat, slightly curved | not hairy | 3 | 5-8.5× 0.4-1 | loose | 2-5 | oblateolate- obovate | 3.5-6.5 | 5 |

Turkey, the NW neighbouring country of Iran, with more than 40 species is one of the most important diversification centers of the genus (McNeill 1967).

During herbarium studies of *Minuartia* material deposited at TARI (abbreviations according to Holmgren & Holmgren 1998), we found a herbarium specimen from Azerbaijan province (NW Iran) which morphologically conformed to the description of neither the species mentioned in Flora Iranica (Rechinger 1988) nor Flora of Turkey (McNeill 1967). More studies displayed that this herbarium material is related to two species i.e. *M. rimarum* (Boiss. & Bal.) Mattf., a narrow endemic to central and southern Turkey, and *M. umbellulifera* (Boiss.) McNeill, known from Iran (Assadi 1984) and Turkey (McNeill 1962). Further studies showed that the material represent a new species; differ from two above species mainly in vegetative characters (Table 1).

MATERIALS AND METHODS

Measurements of the morphological characters were made on the dried herbarium specimens. Pollen grains and ripe seeds were taken from the herbarium material deposited at TARI and were studied using SEM microscopy. Pollen grains and seeds were stabilized on aluminium stocks and coated with a thin layer of gold using coating equipments. Then, the specimens were studied using Scanning Electron Microscope, model LEO 440 at Islamic Azad University, Research and Science Branch. Micro-morphological measurements were performed using Carnoy, a digital measurement software (Schols et al. 2002).

RESULTS AND DISCUSSIONS

Morphology


Minuartia sabalanica Assadi & Mostafavi, sp. nov.

Fig. 1.

Holotypus. Iran, Azerbaijan, Sabalan Mts., 2900 m, 27. Jul. 1972, Foroughi 6120 (TARI).

Species nova differt a *M. umbellulifera* caulibus glabris (nec parte superioribus glanduloso pilosis), foliis linearibus, planis (nec setaceis), 3- nervis (nec uninervis), distantibus (nec densis), petalis oblanceolato-obovatis (nec oblanceolatis), pedicellis 3.5-6.5 mm longis (nec 6-11), seminibus 5 (nec 9-16), a *M. rimarum* caulibus in parte inferioribus glabris (nec glanduloso pilosis), 7.5- 11 cm longis (nec usque ad 8 cm longis), foliis glabris (nec glanduloso pilosis), pedicellis 3.5-6.5 mm longis (nec 5-14.5).

Perennial, 7-11 cm high, glandular-pubescent on the inflorescence. Stems erect. Leaves linear-canaliculate, apiculate, distinctly 3-veined, with narrow membranous margin toward the base. Inflorescence 2-5- flowered,

Fig. 1. *Minuartia sabalanica*.




Fig. 2. Distribution of *Minuartia sabalanica* (●), *M. rimarum* (□) and *M. umbellulifera* (■).

not contracted. Pedicels 3.5-6.5 mm long. Bracts 1.8-3× 0.5-1.1 mm long, ovate-lanceolate to lanceolate, obtuse to apiculate, distinctly 3-veined, with a narrow membranous margin at the base. Sepals 3.5-5×0.6-1.5 mm long; outer sepals slightly longer than the inners, ovate-lanceolate, distinctly 3-veined, with narrow membranous margin toward the base, incurved. Petals much longer than sepals, 5.1-6.1× 1-1.7 mm, oblanceolate-obovate, cuneate at the base, without claw, white. Stamens yellow. Capsule slightly shorter to slightly longer than sepals, 3.8-4.8×1.9-2 mm, cylindrical. Styles 1.2 mm long. Seeds 5 per capsule, 0.9-1×0.6-0.7 mm long, reniform, light brown, verrucate on the surface.

Minuartia sabalanica is morphologically related to *M. umbellulifera*, but differs from it mainly following characters: leaf shape, leaf veins, pedicel length (Tab.1; Mc Neill 1962, 1967; Assadi 1984). The new species is also related to *Minuartia rimarum*, known only from the central and south parts of Turkey differs from it in having following characters which also listed in Table 1: leaf veins, leaf shape (Mc Neill 1967; Rechinger 1988). In addition, a quite long distance is observed between the distribution of *M. sabalanica* and *M. rimarum* (Fig. 2). Considering the obvious geographical distances between *M. rimarum* and *M. sabalanica*, and differences in some diagnostic

characters listed in Table 1, *M. sabalanica* should be mentioned as a distinct species.

Micromorphology

Pollen grain

The results of palynological studies is represented in table 2. Some characters such as, pollen shape, ornamentations, pore number, pore diameter, annulus diameter, were examined in this study (terminology according to Punt & Hoen 1995; Perveen & Qaiser 2006). The observations displayed that pollen grains of both species *M. sabalanica* and also *M. umbellulifera* have rather rounded polyhedral pollen grains (Fig. 3-a,b). Their measures for the quantitative characters plus states of the qualitative characters are listed in Table 2.

The pollen grains in *M. sabalanica* and *M. umbellulifera* are pantoporate with respectively seven and six detectable pores on the equatorial view, and also with scabrate-punctate ornamentations (Table 2, Figs. 3-a,b,c,d).

Seed

The results of seed micro morphology are presented in table 3. Qualitative characters including seed shape, cell shape, cell ornamentation, cell ornament insertion, cell margin shape and also quantitative characters including length and width of seed, length and width of

Table 2. Comparison of palynological data in *Mimuartia sabalanica* and *M. umbellulifera*. Abbreviations used: E: equatorial diameter; P: polar axis length; D: pore diameter (annulus included); R: pore diameter; d: shortest distance between the pores; N: number of pores

| Characters | E(μm) | P(μm) | D(μm) | R(μm) | d(μm) | N(μm) | D/d(μm) | Ornamentation |
|-------------------------|--------|--------|-------|-------|-------|-------|---------|-------------------|
| species | | | | | | | | |
| <i>M. sabalanica</i> | 20.287 | 23.943 | 4.155 | 3.153 | 5.047 | 7 | 0.823 | Scabrate-punctate |
| <i>M. umbellulifera</i> | 23.674 | 26.801 | 5.074 | 4.227 | 5.901 | 6 | 0.859 | Scabrate-punctate |

Table 3. Comparison of seed micromorphological data in *Mimuartia sabalanica* and *M. umbellulifera*. Abbreviations used: SL: seed length; SW: seed width; CL: cell length; CW: cell width; CD: cell distance; SS: seed shape; CS: cell shape CO: cell ornamentation; COI: cell ornament insertion; CMS: cell margin shape.

| Characters | SL(μm) | SW(μm) | SL/SW(μm) | CL(μm) | CW(μm) | CL/CW(μm) | CD(μm) | SS | CO | CMS |
|------------|--------|--------|-----------|--------|--------|-----------|--------|----|----|-----|
| Species | | | | | | | | CS | | |

| | | | | | | | | | | | |
|-------------------------|----------|---------|-------|---------|--------|-------|-------|----------|-----------|-----------|---------|
| <i>M. sabalanica</i> | 1027.347 | 648.33 | 1.584 | 72.593 | 30.694 | 2.365 | 1.821 | reniform | polygonal | verrucate | incised |
| <i>M. umbellulifera</i> | 1062.086 | 905.228 | 1.172 | 111.374 | 34.53 | 2.225 | 2.808 | reniform | oblong | verrucate | incised |

cell and distance between cells, were examined (terminology according to Bittrich 1993).

Minuartia sabalanica has an almost reniform seed shape covered with irregular polygonal cells (Fig. 1-k, Fig. 4-a, c). Cell shape is a diagnostic character for distinguishing *M. sabalanica* from *M. umbellulifera* (Table 3, Fig. 4- c,d). Cell margin shape in *M. umbellulifera* is quite incised (Fig. 4-d).

The results indicate that seed micromorphology provides more effective characters for distinguishing these two species from each other.

It should be mentioned that the species *M. rimarum* does not occur in Iran. Therefore, *M. umbellulifera*, the closest species found in Iran, was studied micro-morphologically.

ACKNOWLEDGMENT

The authors wish to thank Mr. M. Mehranfard the artist of Plant Pests and Diseases Research Institute for the drawing of Fig. 1, and Mrs. S. Eshghi, for preparing SEM micrographs.

REFERENCES

- Assadi, M. 1984: New species and new plant records from Iran. - Iran. J. Bot. 2 (2):83-93.
- Bittrich, V. 1993: Caryophyllaceae. In: Kubitzki K., Rohwer J. G. and Bittrich V. (eds.), The Families and Genera of Vascular Plants, vol. 2, Flowering Plants Dicotyledons: Magnoliid, Hamamelid and Caryophyllid Families. -Springer, Germany, 206-236.
- Boissier, E. 1867: Alsine in Flora Orientalis, vol. 1: 669-704. -Basileae & Genevae.
- Halliday, G. 1964: Minuartia in Tutin, T. G., Heywood, V. H., Burges, N. A. & Web, D. A. (eds.). Flora Europaea: 1: 125-132. - Cambridge at the University Press.
- Holmgren, P. K. & Holmgren, N. H. 1998: (continuously updated): Index Herbariorum (online at <http://sciweb.nybg.org/science2/IndexHerbariorum.asp>).
- Kamari, G. 1997: Minuartia L. in Strid, A. & Tan, K. (eds.). Flora Hellenica.- vol. 1: 170-191. – Koeltz Scientific Books, Germany.
- Mc Neill, J. 1962: Taxonomic studies in the Alsinoideae: I. Genetic and infrageneric groups. - Notes. R. G. B. Edinburgh 24: 79-155.
- Mc Neill, J. 1963: Taxonomic studies in the Alsinoideae: II. A revision of the species in the orient. -Notes. R. G. B. Edinburgh 24:241-404.
- Mc Neill, J. 1967: Minuartia in P. H. Davis, Flora of Turkey and the East Aegean Islands. vol. 2: 38-67. – Edinburgh at the University Press, Edinburgh.
- Meikle, R. D. 1977: Minuartia in Flora of Cyprus. vol. 1: 265-273. -Bentham-Moxon Trust Royal Botanic Gardens, Kew.
- Parsa, A. 1951: Minuartia in Flore de'l Iran. - vol. 1: 1159-1177. -Tehran.
- Perveen, A. & Qaiser, M. 2006: Pollen flora of Pakistan-LI-Caryophyllaceae. Pak. J. Bot. 38 (4): 901-915.
- Punt, W. & Hoen, P. P. 1995: Caryophyllaceae in The Northwest European Pollen Flora. Review of Palaeobotany and Palynology. VII. 88 (1-4): 83-272. -Elsevier, Amsterdam.
- Rechinger, K. H. 1988: Minuartia in K. H. Rechinger, Flora Iranica. no. 163: 28-53. -Academische Druck & verlagsanstalt Graz.
- Schischkin, V. L. 1936: Minuartia in Flora of the U.S.S.R., vol. 6: 482- 516. - Translated by (Smithsonian Institution and the National Science Foundation, Washington DC. Program for Scientific Translation, Jerusalem, Israel).
- Scholes, P., Dessein, S., D'Hondt, C., Huysmans, S. & Smets, E. 2002: CARNOY: a new digital measurement tool for palynology.- Grana vol. 41: 124-126.




Fig. 3. SEM micrographs of pollen morphology of *Minuartia sabalanica*: a) general view c) pore and ornate-ments. scales: a = 3 μ m b = 1 μ m. *M. umbellulifera*: b) general view d) pore and ornate-ments. scales: a= 3 μ m. b=1 μ m.




Fig. 4. SEM micrographs of seed morphology: *Minuartia sabalanica*: a) general view c) testa cells. scales: a= 100 μ m. c= 10 μ m. *Minuartia umbellulifera*: b) general view d) testa cells. scales: a= 100 μ m. d= 2 μ m.

