

Original Article

Antioxidant Compounds and Activity of the Medicinal Plants *Tagetes* sp. and *Clinopodium* sp. from Indigenous Communities in Mexico**Running title:** Antioxidant compounds and activity of *Tagetes* sp. and *Clinopodium* sp.**Mónica L. Pérez-Ochoa, Araceli M. Vera-Guzmán and José Luis Chávez-Servia***

CIIDIR-Oaxaca, Instituto Politécnico Nacional, Santa Cruz Xoxocotlán, Oaxaca, México

Article History

Received: 03 March 2022
Accepted: 26 August 2023
© 2012 Iranian Society of
Medicinal Plants.
All rights reserved.

Keywords

Spectrophotometry
Traditional knowledge
Phenolic compounds
Gastrointestinal disorders

***Corresponding author**
jchavez@ipn.mx

ABSTRACT

Knowledge about the therapeutic properties and efficacy conferred by several plants to indigenous communities is based on observations, traditional knowledge, and forms of use or application within local, cultural, and territorial contexts. However, to some extent, favorable responses to these plants depend on the form in which a plant is used and its phytochemical composition. This work aimed to evaluate the contents of total polyphenols and flavonoids and antioxidant activity and determine qualitative phytochemical profiles of *T. lucida*, *T. lunulata*, *C. mexicanum* and *C. macrostemum* from samples of plants collected in four indigenous regions of Oaxaca, Mexico. The results showed that the contents of phenolic compounds and antioxidant activity differed significantly among the evaluated species and that the effect of species was greater than the ecological-environmental effects caused by the origins of the samples. Tannins, alkaloids, steroids, terpenoids, and coumarins were qualitatively identified in all evaluated species, and emodins were detected only in *T. lucida*. Among the communities visited, it was documented that all the species are used to treat gastrointestinal disorders, and antimicrobial, calming and anti-inflammatory properties were recorded for all four species. The effectiveness conferred to these plant species in the communities to treat ailments, including endemic diseases, is partly due to the complex phytochemical composition and the beneficial properties released during infusion or decoction.

INTRODUCTION

The use of medicinal plants transcends territorial, cultural and temporal barriers in such a way that treatment with these plants is now prevalent [1]. For example, in genomic medicine, it is possible to evaluate natural extracts of fruits or plants by gene expression of antioxidant metabolic enzymes using biological models [2] to define potentialities in medicinal plants. The World Health Organization (WHO) reports that 80% or more of the countries in the world use herbal medicine as part of traditional or complementary treatments, with greater prevalence in Africa, America, some European countries and the Eastern Pacific region due to the available plants and indigenous knowledge in these areas [3]. The WHO has indicated that there is not enough information on the potential of these plants for pharmacological use and their phytochemical compositions and that research on the safety, monitoring, and efficacy of

commonly used plants, such as those that are incorporated into herbal medicines or their derived products that are taken daily, is lacking [2-3]. For example, *Prosopis juliflora* (Sw) DC. (mezquite in Spanish) has potential antiproliferative effects against human prostate LNCaP cells [5], but more pharmacological studies are necessary. Zhang [6] and Lee et al. [7] pointed out that the products, practices, and practitioners of traditional and integrative medicine are part of the first medical aid provided by all societies.

Among indigenous communities in Mexico, species of the genera *Tagetes* (Asteraceae) and *Clinopodium* (Labiatae) are widely used in rituals and ethnoreligious ceremonies (e.g., Day of the Dead, family festivals or social events) and traditional medicine [8-11]. Leaves, flowers, and stems of *T. lucida* Cav. and *T. lunulata* Ortega, also known as 'cempoalxóchitls' (marigolds), are prepared in the

form of infusions or decoctions in indigenous communities to treat gastrointestinal conditions, menstrual pain, cough, and endemic diseases locally called 'empacho', 'mal de ojo' and 'susto' (ethnodiseases) and as a relaxant [12-14]. The aerial parts of the 'poleos' (pennyroyal), *Clinopodium macrostemum* (Moc. & Sessé ex Benth.) Kuntze (Syn. *Satureja macrostema*) and *C. mexicanum* (Benth.) Govaerts are also used to treat stomach pain, intestinal cramps, indigestion, and gastrointestinal infections and to reduce the effects of excess alcohol consumption [10, 15, 16].

Previous studies have revealed anti-inflammatory, antibacterial and antiulcer properties and antioxidant activities in extracts of different species of *Tagetes* and *Clinopodium* [17-19]. For example, in *T. lucida*, glycosylated flavonols, phenolic acids, aromatic acids and hydroxylated coumarins have been identified, compounds that confer high antioxidant activity and produce antidepressant, anxiolytic, sedative, antifungal, and antibacterial effects, protecting against bacteria that cause intestinal infections (e.g., *E. coli*, *Salmonella* sp. and *V. cholerae*); these compounds also produce analgesic effects and help prevent diabetes and obesity [17, 20-23]. In addition, *C. macrostemum* extracts are hepatoprotective, renoprotective, and antidiarrheal and relieve intestinal pain [24-27]. Analgesic and sedative effects have been reported in *C. mexicanum* [28]. All these findings indicate that these medicinal plants have varying phytochemical compositions that need to be understood and that the ecological conditions in which the plants grow lead to variations in plant composition and compound concentration. In this context, the objective of this study was to evaluate the contents of total phenols and flavonoids and antioxidant activity and to determine a qualitative phytochemical profile of *T. lucida*, *T. lunulata*, *C. mexicanum* and *C. macrostemum* from plant samples collected in four indigenous regions of Oaxaca, Mexico.

MATERIALS AND METHODS

Documentation of Communities and Collection of Plant Material

At the flowering stage, samples of *T. lucida* Cav., *T. lunulata* Ortega, *C. macrostemum* (Moc. & Sessé ex Benth.) Kuntze and *C. mexicanum* (Benth.) Govaerts (syn.: *Satureja oaxacana*; *Satureja mexicana*) were collected in two locations for each species in the

Mixteca, Valles Centrales, Sierra Sur, and Sierra Norte regions from Oaxaca, Mexico. Sample collection for phytochemical analysis was performed at two locations per species from October to December 2018 in regions with different climatic conditions. In each community or locality, ecogeographic descriptors of the soil and climate were recorded based on geodetic maps provided by the National Institute of Statistics and Geography (INEGI) [29]. Furthermore, the altitude, latitude and longitude of the sites were recorded by a GPS unit. In addition, ethnobotanical information on the uses of each species was obtained through rapid surveys with key members in eight communities of the regions visited (Table 1).

Preparation of Samples for Laboratory Analysis

The plant material was washed with running water and rinsed with distilled water, left to dry in the shade at room temperature, and subsequently ground in an electric mill (Krupps®, model GX4100, Mexico) for 3 min (at intervals of 1.5 min) until a homogeneous powder with a particle size of 0.6 mm was obtained. The dry milled samples were stored in amber flasks at 4 °C until analysis. The determination of total polyphenols and flavonoids was made based on 0.1 g of ground sample, which was mixed with 25 mL of 80% ethanol. The evaluation of antioxidant activity (DPPH and FRAP) was performed with a 0.1 g sample mixed with 25 mL of 80% methanol. In both cases, the mixtures were homogenized (Ultra Turrax T25 Digital, IKA, Germany) for 60 s (in 30 s intervals) and centrifuged at 11500 rpm (refrigerated Eppendorf 5810R centrifuge) for 15 min at 10 °C, and the supernatant was extracted for the corresponding determinations, in each case with three replicates per extract.

Evaluation of Phenolic Compounds and Antioxidant Activity

Total Polyphenols

The evaluation of polyphenols was performed based on the Folin-Ciocalteu method [31], which is used to measure the concentration of total phenolic compounds based on the transfer of electrons from the extract to phosphomolybdic acid/phosphatidic acid complexes in an alkaline medium, and a concentration-dependent blue color was obtained. The absorption was measured at 750 nm using a spectrophotometer (UV-1800 Shimadzu Corporation

Kyoto, Japan). The concentration of each sample was obtained with reference to a calibration curve of the standard gallic acid (0.021-0.165 mg/mL), and the results were expressed as milligrams of gallic acid equivalents per gram of dry weight (mg GAE/g dw).

Flavonoids

The estimation of flavonoids was made using two complementary colorimetric methods based on the integration of complex compounds of different chemical natures. The first method was based on the reaction of aluminum chloride (AlCl₃) with flavonoids [32], and it tests specifically for flavonols and luteolin (flavone). In this case, the absorbance values were recorded at 410 nm using a spectrophotometer (Shimadzu UV-1800, Kyoto, Japan), and the concentration was expressed as mg equivalents of quercetin per gram of dry weight (mg QE/g dw) with reference to a quercetin calibration curve (0.01-0.07 mg/mL). The second method was

performed with sodium nitrite (NaNO₂) in an alkaline medium [33] and was used to test for catechins [34]. Spectrophotometric evaluations were performed at 510 nm according to the reference curve of catechin (0.01-0.12 mg/mL), and the results were expressed in mg catechin equivalents per gram of dry weight (mg CE/g dw).

Antioxidant Activity Determined by the DPPH and FRAP Methods

The DPPH (2,2-diphenyl-1-picrylhydrazyl, Sigma-Aldrich) antioxidant activity was evaluated by the method described by Brand-Williams [35]; this method involved recording absorbances at 517 nm in a spectrophotometer (Shimadzu UV-1800, Kyoto, Japan) and calculating the concentration based on the standard Trolox curve (6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid) (0.13-1.19 µmol/mL).

Table 1 Species collected belonging to *Tagetes* and *Clinopodium* and ecological descriptions of the communities and municipalities sampled in Oaxaca, Mexico.

Species and communities at the sampling sites	Altitude (m), latitude (N), longitude (W)	Predominant climate in the communities ^a	Temperature (°C) ^a and average rainfall (mm)	Predominant soil types ^b
<i>T. lucida:</i>				
Santa Lucía Miahuatlán	1966 16° 11' 0" 96° 38' 0"	Temperate subhumid with rainfall in summer (72.4%)	12-22 700-2000	Leptosol, luvisol and regosol
San Pedro Cajonos	2234 17° 10' 13.6" 96° 18' 10.2"	Temperate subhumid with rainfall in summer (90.3%)	16-20 1200-2000	Luvisol
<i>T. lunulata:</i>				
San Pablo Huitzo	1813 17° 17' 4.3" 96° 57' 2.8"	Temperate subhumid (57.45%) and semiwarm subhumid (41.99%) with rainfall in summer	14-22 600-1000	Regosol, leptosol, and vertisol
Tamazulapan del Espíritu Santo	2067 17° 03' 30.1" 96° 03' 44.6"	Temperate subhumid with rainfall in summer (84.03%)	14-20 800-1500	Acrisol, luvisol, and cambisol
<i>C. macrostemum:</i>				
Santa Catarina Lachatao	2831 17° 07' 5.2" 96° 31' 41.8"	Temperate subhumid with rainfall in summer (95.84%)	10-20 800-1200	Acrisol and cambisol
San Pedro y San Pablo Ayutla	2261 17° 00' 45" 96° 07' 33"	Semiwarm subhumid (36.55%), temperate subhumid (28.26%), and temperate wet (24.09%) with rainfall in summer	14-24 700-1500	Acrisol, cambisol, and luvisol
<i>C. mexicanum:</i>				
Santo Domingo Yanhuitlán	2236 17° 33' 1.7" 97° 22' 17.8"	Temperate subhumid (81.04%) with rainfall in summer and other seasons	14-18 600-800	Leptosol, vertisol, and phaeozem
San Miguel Tixá	2274 17° 31' 20.9" 97° 30' 56.0"	Temperate subhumid (93.33%) with rainfall in summer and other seasons	14-20 700-900	Leptosol, luvisol, and vertisol

^aINEGI [29]; ^bSoil type based on the IUSS Working Group WRB classification [30].

The antioxidant activity was also evaluated by the FRAP method (iron-reducing power) with reference to the approach proposed by Benzie and Strain [36]. Absorbance was recorded at a wavelength of 593 nm, and quantification was performed based on the Trolox calibration curve (0.05-1.0 $\mu\text{mol/mL}$). The results of both determinations were expressed as micromolar equivalents of Trolox per gram of dry weight ($\mu\text{mol TE/g dw}$).

Standard Qualitative Determination of Phytochemical Compounds

Qualitative phytochemical evaluation of the extracts was carried out by standard methods based on changes in color or the formation of precipitates, which were recorded as present (+/1) or absent (-/0). This exploratory procedure is frequently used in studies of plant extracts to determine the presence of groups of bioactive compounds in plants [37]. In this work, 7 g of each powder sample was mixed with 70 mL of 80% methanol and homogenized for 60 s (Ultra Turrax T25 Digital, IKA, Germany). The resulting extract was vacuum filtered, and liquid-liquid extraction was successively carried out with solvents of increasing polarity from hexane and chloroform to ethyl acetate, recovering the phases with each solvent. The final fractions were stored at 4 °C until evaluation.

Qualitative phytochemical analysis was performed using the methodologies reported by Savithamma [37] and Marcus [38]. Tannins: drops of 1% lead acetate were added to 1 mL of the extract, and the presence of tannins was recorded when a yellow precipitate was observed. Saponins: 1 mL of the extract was mixed with 2 mL of distilled water and stirred vigorously for 2 min; the presence of the compounds was recorded following foam formation in the solution. Terpenoids: First, 1.5 mL of the extract was mixed with 1 mL of chloroform. Then, 1 mL of concentrated H_2SO_4 was added, and the presence of terpenoids was indicated by the formation of a reddish-brown ring. Steroids: Three drops of concentrated H_2SO_4 were added to 3 mL of the extract, and a red coloration indicated the presence of steroids. Alkaloids: Three drops of 1% HCl were added to 1.5 mL of extract, and the solution was mixed; then, two drops of Wagner's reagent (2 g of I_2 and 6 g of KI dissolved in 100 mL of distilled water) were added to the solution, and the formation of a reddish-brown precipitate indicated the presence of alkaloids. Flavonoids: One or two drops of 1 M NaOH solution

was added to 1 mL of extract, causing the solution to turn an intense yellow color. Then, two to three drops of 1 M HCl were added, and when the solution became colorless, it indicated the presence of flavonoids. Coumarins: 1 mL of 10% NaOH was added to 1 mL of the extract, and a yellow hue indicated the presence of coumarins.

Statistical Analysis

Based on the data generated in the quantitative analysis of phenolic compounds and antioxidant activity, analysis of variance was performed using a random linear model, where the origins of the samples collected from each species were assumed to be nested by species. To evaluate the differences between species and sample origins, multiple comparisons of means were made with the Tukey test ($P < 0.05$). In addition, a Pearson correlation analysis ($P \leq 0.05$) was performed between average values of total polyphenols, flavonoids and antioxidant activity. For the qualitative determinations, descriptive analyses were performed. Statistical analyses were performed with the SAS statistical package [39].

RESULTS AND DISCUSSION

Community uses of *Tagetes sp.* and *Clinopodium sp.*

The indigenous communities of Oaxaca, Mexico, widely use marigolds (*Tagetes sp.*) and 'poleos' (*C. sp.*) for ornamental, medicinal, ritual and nutritional purposes, and the plants are a part of their cultures and traditions (Table 2). This finding coincides with previous reports regarding the use of these species [10, 27]. In the study region, it was observed that among Zapotec communities (in the subtropical or warm zone) of the Central Valleys and Sierra Norte of Oaxaca, Mexico, *C. macrostemum* was the main medicinal plant. In addition, in the temperate to humid temperate zones occupied by the Mixtec group, *C. mexicanum* was used exclusively. *T. lucida* (pericón) has more medicinal functions. In contrast, *T. lunulata* has more ritual and religious uses, e.g., as an altar decoration in honor of the dead in a traditional Mexican ceremony celebrated on November 1 and 2. Subsequently, the plants are taken to cemeteries, where they are used to decorate tombs. Both *T. lucida* and *T. lunulata* have a variety of medicinal uses and are distributed in temperate zones (Table 2).

Table 2 Descriptions of traditional uses of *Tagetes* and *Clinopodium* species among indigenous communities from Oaxaca, Mexico.

Species sampled	Local names in Spanish and native languages	Examples of ethnobotanical uses ¹
<i>T. lucida</i>	Pericón, <i>tiex</i> in Zapoteco (native language)	Food: The plant is added to boiled corn for flavor. An infusion of the sweetened aerial part of the plant is consumed. Ornamental and ceremonial: The plant is used as an adornment for the image of the Virgin Mary in the Catholic religion. Medicinal: An infusion of the aerial part of the plant relieves stomach pain, diarrhea, 'empacho', irregular heartbeat, stomach discomfort, menstrual pain and is used as an anti-inflammatory; the aerial part of the plant is soaked in 'aguardiente' (sugarcane alcohol) relieve stomach pain. The boiled root is used to treat diabetes and cancer.
<i>T. lunulata</i>	Cempasúchil cimarrón, flor de todos santos, flor de muerto chiquita o cempasúchil silvestre; flor de angelito, cempasúchil, cempasúchil hembra; flor de muerto; cempasúchil chiquito; flor de niño (native language).	Ritual and ceremonial: The plant is used as an adornment of the altar of the dead. Ornamental: The plant is appreciated for its aroma and the showy color of its petals. Medicinal: An infusion of the aerial part of the plant relieves stomach pain.
<i>C. macrostemum</i>	Poleo, hierba de borracho, rosa de borracho and <i>Gú guázz</i> , <i>guiezza</i> , <i>quieutzu</i> in Zapoteco (native language).	Food: The leaves are consumed cooked or added during the cooking of beans (<i>Phaseolus</i> sp.) and tamales and served as a condiment. An infusion of the aerial part of the plant is consumed in water, or the leaves and stems are steeped in 'aguardiente' (sugarcane alcohol) or mezcal for consumption as an alcoholic beverage. Ritual and ceremonial: It is given to guests at mayordomías, weddings and Christmas parties. Medicinal: Relieves stomach pain, diarrhea, heartburn, fever, symptoms of excess alcohol intake, and high blood pressure.
<i>C. mexicanum</i>	Chepito, hierba del borracho, toronjil, poleo, hierba de menta and <i>Gú guázz</i> in Zapoteco (native language)	Food: An infusion of the sweetened aerial part of the plant is consumed. Medicinal: An infusion or maceration in brandy relieves stomach pain caused by excess alcohol intake. The plant is also an analgesic and sedative and can induce sleep.

¹A synthesis of ethnobotanical information gathered during a quick survey of species uses among indigenous communities from Oaxaca, Mexico, complemented with information provided by previous works [10,28, 40-45].

Table 3 Significance of the square means from the analysis of variance of total polyphenol and flavonoid contents and antioxidant activity in two species of *Tagetes* and two species of *Clinopodium*.

Source of variation	Total polyphenols	Flavonoid equivalents		Antioxidant activity	
		Quercetin	Catechin	DPPH	FRAP
Species	7080 ** (98.7) ¹	7236 ** (99.7)	18268 ** (94.6)	1096476 ** (99.5)	1740662 ** (99.3)
Community of origin of samples within each species	86.8 ** (1.2)	16.3 ** (0.2)	1030.3 ** (5.3)	4029.1 ** (0.4)	10445.2 ** (0.6)
Error	2.4 (0.1)	1.5 (0.1)	7.9 (0.1)	1042.1 (0.1)	273.5 (0.1)
C.V. (%)	3.6	6.1	4.1	8.9	3.7

**Significant at $P \leq 0.01$; ¹values in parenthesis indicate percentage of variance explained by the total variance estimated in the linear model; C.V. = coefficient of variation.

The fresh and dry stems, leaves and flowers of *C. macrostemum*, *C. mexicanum*, *T. lucida* and *T. lunulata* are widely used in medicinal infusions and boiled solutions to counteract stomach and gastrointestinal discomfort (Table 2). According to

the interviewees, *C. macrostemum* and *C. mexicanum* are preferably consumed in infusions or decoctions and served as hot drinks for the treatment of diarrhea, stomach pain caused by ingestion of food or alcoholic beverages and menstrual pain; these findings coincide with reports by UNAM [16] and Arrazola-

Guendulay [10] regarding the curative uses of these plants for stomach pain, nausea, indigestion, fever, diarrhea, and intestinal cramps (including the symptoms of ethno-diseases) and to alleviate symptoms of excessive alcohol consumption. In addition, these species are used as condiments in the preparation of food; combined with ground beans (locally called 'pasta'), tamales (local dishes), or refreshing drinks; or used in the preparation of soups and vegetables similar to the so-called *quelites*, edible leaves and tender stems of the genera *Amaranthus*, *Portulaca*, *Chenopodium* and others. In addition, these species are included in the preparation of a liqueur: the leaves and stems, preferably green, are added to equal volumes of 0.25, 0.5 or 1.0 L of 'aguardiente' (alcohol of sugarcane, *Saccharum officinarum* L.) or mezcal distilled from *Agave angustifolia* Haw. That is, *Clinopodium* species have diverse uses among indigenous communities. Similar aspects were documented for *T. lucida* and *T. lunulata*, which can be prepared similarly and are used to treat stomach pain, diarrhea, menstrual pain, fever, vomiting and ethno-diseases such as 'empacho', 'mal de ojo' and 'susto' [10, 12-14].

Phenolic Compounds and Antioxidant Activity

Based on the analysis of variance, significant differences ($P \leq 0.01$) were found between the species evaluated (*T. lucida*, *T. lunulata*, *C. macrostemum* and *C. mexicanum*) and between the communities of origin of the samples of each species in terms of the contents of total polyphenols and flavonoids (quercetin and catechin) and antioxidant activity evaluated by the DPPH and FRAP methods. It is important to note that the variation attributed to the effects of the growth environment of the sampled plants was significantly lower (0.2 to 5.3%) than the variation estimated for the effect of species (94.6 to 99.3%); these findings are based on the linear model used to perform the analysis of variance (Table 3) and coincided with the results reported by Cicevan [46], who did not find a consistent difference in the contents of phenols and flavonoids between species and varieties of *Tagetes* subjected to water stress. Consequently, these results indicate that the variation in the contents of phenolic compounds and antioxidant activity between samples within the same species can be used to determine the species composition, as has been reported in other medicinal plants [47, 48].

Polyphenols and flavonoids in medicinal plants play an important communitarian role in the prevention and treatment of ethnodiseases and metabolic disorders of human health. As part of the potential application in medicinal chemistry, it is important to elucidate such a mechanism; for example, Ullah et al. [49] found that flavonoids are associated with the prevention of cancer and anti-inflammatory, antiviral, neuroprotective and cardio-preventive effects. The biological activities depend on the specific flavonoid, mode of action, bioavailability and ease of release from plant extracts. In this sense, Nuñez et al. [50] tested whether *T. erecta* L. extracts have antidiabetic and anti-obesity properties using *Caenorhabditis elegans* as a biological model.

Species of *Tagetes* and *Clinopodium* are used by indigenous communities in Oaxaca, Mexico, for the treatment of gastrointestinal disorders (Table 2), and the healing effects of these plants are directly based on their phytochemical composition. For example, the phenol and flavonoid contents of these species are associated with antimicrobial and anti-inflammatory properties and relaxing effects [27,28,51,52]. The results show that the contents of polyphenols and flavonoids in *Tagetes* ranged from 29.6 to 71.7 mg GAE/g and from 17.8 to 95.6 mg QE/CE/g, respectively; in *Clinopodium*, the variation ranged from 30.1 to 39.5 mg GAE/g and from 6.3 to 96.1 mg QE/CE/g. *T. lucida* had higher contents of total polyphenols and flavonoids than *T. lunulata*, a pattern that is reflected in greater antioxidant activity; similar patterns were also recorded in *C. mexicanum* with respect to *C. macrostemum* (Table 4). The determination of general or specific compounds in medicinal plants guides future studies or specific pharmacological tests. For example, in traditional Chinese medicine, *C. chinense* is reported to be a therapeutic option for cardiac fibrosis due to first reports of an antifibrosis mechanism in rat models with myocardial injury [53]. However, more evaluations are necessary because not always all species of the same genera confer similar effects; for example, Morales-Torres *et al.* [54] determined the absence of nephroprotective effects in the evaluation of aqueous extracts of *C. vimineum* (L.) Kuntzie using a renal disorder model with Wistar rats.

The variation in phenolic compounds and antioxidant activity between samples collected in different communities reflected different patterns between species. For example, the contents of total

polyphenols and quercetin-equivalent flavonoids and antioxidant activity differed between *T. lucida* samples according to community origin; the values of these measures were also significantly higher in *T. lucida* than in *T. lunulata*, *C. macrostemum* and *C. mexicanum*. Therefore, to a great extent, the differences occur at the species level and are not due to the different ecological origins of the samples, although it was necessary to obtain different samples to estimate the variation in composition within each sample. Within each species, there were significant differences in the total polyphenol content according to the sample origin, but there were no differences in the contents of quercetin-equivalent flavonoids within the species *T. lunulata*, *C. macrostemum* and *C. mexicanum*. The antioxidant activity in the

samples of *T. lucida* and *C. mexicanum* differed significantly according to one or both evaluation methods (DPPH or FRAP), but this result was not observed for *T. lunulata* or *C. macrostemum* (Table 5). This indicates that the growth environment of the plant has an influence on its contents of phenolic compounds and antioxidant activity but that the pattern changes from species to species. These patterns have some similarities to those reported by Dunkić [55], who differentiated *C. dalmaticum*, *C. pulegium*, *C. serpyllifolium* and *C. thymifolium* by the contents of tannins, phenolic acids, flavonoids, and total polyphenols, and to those reported by Cicevan [46] and Li *et al.* [56], who established differences between *Tagetes erecta*, *T. patula* and *T. tenuifolia* according to total phenol and flavonoid contents.

Table 4 Differences in the contents of phenolic compounds and antioxidant activity among *Tagetes* and *Clinopodium* species collected in Oaxaca, Mexico.

Species evaluated	Total polyphenol content (mg GAE/g)	Flavonoid equivalent		Antioxidant activity	
		Quercetin (mg QE/g)	Catechin (mg CE/g)	DPPH (μmol TE/g)	FRAP (μmol TE/g)
<i>T. lucida</i>	71.7 a ¹	48.9 a	95.6 a	725.8 a	901.7 a
<i>T. lunulata</i>	29.6 c	17.8 b	32.9 c	257.2 b	264.5 c
<i>C. macrostemum</i>	30.1 c	6.3 c	51.7 b	179.9 c	233.4 d
<i>C. mexicanum</i>	39.5 b	6.7 c	96.1 a	282.9 b	380.6 b

¹In columns, means with the same letter are not significantly different (Tukey's test, $P \leq 0.05$).

Table 5 Variation in the contents of phenolic compounds and antioxidant activity within species of *Tagetes* and *Clinopodium* collected in Oaxaca, Mexico, according to the community of origin of the samples.

Species and community of origin of the samples	Total polyphenols (mg GAE/g)	Flavonoid equivalent		Antioxidant activity	
		Quercetin (mg QE/g)	Catechin (mg CE/g)	DPPH (μmol TE/g)	FRAP (μmol TE/g)
<i>T. lucida</i>					
Santa Lucia Miahuatlán	68.4 b ¹	50.5 a	92.9 c	739.7 a	878.6 b
San Pedro Cajonos	74.9 a	47.4 b	98.2 b	711.9 a	924.9 a
<i>T. lunulata</i>					
San Pablo Huitzo	31.0 e	18.6 c	31.5 g	264.7 bc	276.2 e
Tamazulapám del Espíritu Santo	28.3 f	16.9 c	34.2 g	249.7 c	252.8 ef
<i>C. macrostemum</i>					
Santa Catarina Lachatao	28.7 f	5.6 d	48.7 f	171.6 d	229.7 f
San Pedro y San Pablo Ayutla	31.4 e	7.0 d	54.6 e	188.1 d	237.2 f
<i>C. mexicanum</i>					
Santo Domingo Yanhuitlán	41.7 c	6.9 d	110.6 a	307.0 b	421.0 c
San Miguel Tixa	37.2 d	6.5 d	81.5 d	258.9 c	340.1 d

¹In columns, means with the same letter are not significantly different (Tukey's test, $P \leq 0.05$).

In twelve varieties of *T. patula*, *T. tenuifolia* and *T. erecta*, Cicevan [46] estimated that the total phenol content ranged from 2.52 to 8.38 mg GAE/g and that the total flavonoid content ranged from 2.72 to 7.88 mg CE/g; in this study, the observed variations were slightly higher, with the ranges for *T. lucida* and *T. lunulata* ranging from 28.3 to 68.4 mg GAE/g and

from 31.5 to 98.2 mg CE/g, respectively. This finding indicates that despite the methodological differences between laboratories, there are higher concentrations of these compounds in native Mexican species than in the species evaluated by Cicevan [46], and this finding may also partially explain the beneficial

effects conferred by these species in traditional medicine.

The correlation analysis showed that the contents of total polyphenols were positively and significantly correlated with the antioxidant activity estimated by FRAP ($r = 0.99^{**}$, $P \leq 0.01$) and DPPH ($r = 0.96^{**}$, $P \leq 0.01$) methods, and similar patterns were observed for the contents of flavonoids evaluated according to the quercetin standard (QE) with the DPPH ($r = 0.95^{**}$, $P \leq 0.01$) and FRAP ($r = 0.92^{**}$, $P \leq 0.01$) methods and the catechin standard (CE) with the FRAP ($r = 0.68^{**}$, $P \leq 0.01$) and DPPH ($r = 0.59^{**}$, $P \leq 0.01$) methods. The results show a close association of polyphenols and flavonoids with antioxidant activity, but antioxidant effects of other compounds are not ruled out, as indicated by the findings of Cicevan [46] in species of *Tagetes* and Dunkić [55] in species of *Clinopodium*.

The differential polarity of each solvent used (hexane, chloroform and ethyl acetate) helped to qualitatively determine the presence of different phytochemical compounds. Hexane allowed us to determine the presence of saponins, steroidal terpenoids and alkaloids in *T. lucida* and *T. lunulata*, but the same solvent only helped to determine the presence of alkaloids in *C. mexicanum* and *C. macrostemum*. Specifically, alkaloids were present in all the species evaluated when hexane and ethyl acetate were used but not when chloroform was used, and this result may guide future studies to evaluate the specific concentrations of this group of compounds. Similarly, the results show that the use of chloroform and ethyl acetate as extraction solvents is recommended for the evaluation of the contents of tannins, terpenoids, coumarins and flavonoids in both *Tagetes* and *Clinopodium*. However, the presence of saponins was detected in *C. mexicanum*, *C. macrostemum* and *T. lucida* only with chloroform. Emodins were detected only in *T. lucida* with ethyl acetate (Table 6). Therefore, between and within the species evaluated, the same group of phytochemical compounds was not always observed in samples from two geographical origins. For example, saponins and steroids in *T. lunulata* were detected with hexane, while saponins and emodins in *T. lucida* were detected with chloroform and ethyl acetate; these compounds were not observed with this method in *T. lunulata*, and in this species, steroids were observed using ethyl acetate. These findings indicate that the

samples collected probably did not have a sufficient concentration of compounds to be detected by the qualitative techniques used. However, the use of ethyl acetate and chloroform as extraction solvents shows reactive potential for quantitative determinations of tannins, saponins, terpenoids, steroids, alkaloids, coumarins and flavonoids.

The quantification of phenolic compounds and flavonoids, qualitative determination of compounds (tannins, saponins, terpenoids, steroids, alkaloids, coumarins and emodins), and high antioxidant activity indicate that the evaluated extracts are complex matrices of compounds; this finding partially explains the use of *T. lucida*, *T. lunulata*, *C. macrostemum*, and *C. mexicanum* in the traditional medicine of the indigenous communities of Oaxaca, Mexico. For example, flavonoids in *T. lucida*, including quercetagenin 7-O- β -D-glucoside, quercetagenin-7-O- β -D-glucopyranoside, 6-hydroxykaempferol-7-O- β -D-glucopyranoside and 5,7,4'-trimethoxyflavone, which have high antioxidant capacities, are analgesic and inhibit the growth of enterobacteria and pathogenic fungi [20, 21,23,51,57]. Coumarins (7,8-dihydroxycoumarin, 7,8-dihydroxy-6-methoxycoumarin, and 7-isoprenyloxycoumarin) have also been detected in *T. lucida*. 6,7-Dimethoxycoumarin, herniarin and 6,7,8-trimethoxy-coumarin have anti-inflammatory, analgesic, vasorelaxant, anxiolytic and sedative properties [11,20,48,51,58]. All these findings support the possible medicinal effects conferred on these plants by the indigenous communities of Mexico.

Terpenes confer antimicrobial properties to the plants that possess them. In this study, terpenes were present in the chloroform and ethyl acetate fractions in all the species evaluated. Villa-Ruano [59] identified the terpenes menthone and piperitone oxide in *C. macrostemum*, and the results of the study partially explained the medicinal use of this plant in Oaxaca, Mexico, for stomach pain. The same effect has been described for *C. mexicanum*; the steroid saponins β -sitosterol and daucosterol, which have antimicrobial potential, have been detected in this species [27]. Additionally, it was determined that *T. patula* and *T. lunulata* contain alkaloids (Table 6), compounds associated with anesthetic, stimulant, antimicrobial and anti-inflammatory properties.

Table 6 Screening of phytochemical compounds based on methanolic extracts of hexane, chloroform, and ethyl acetate solvents in *Tagetes* and *Clinopodium* species from Oaxaca, Mexico.

Phytochemical compounds	<i>C. mexicanum</i>		<i>C. macrostemum</i>		<i>T. lucida</i>		<i>T. lunulata</i>		
	Santo Domingo Yanhuitlán	San Miguel Tixá	Santa Catarina Lachatao	San P. y San P. Ayutla	Santa Lucía Miahuatlán	San Pedro Cajonos	San Pablo Huitzo	Tamazulapam del Espíritu Santo	
Hexane									
Tannins	0	0	0	0	0	0	0	0	
Saponins	0	0	0	0	0	1	0	1	
Terpenoids	0	0	0	0	1	1	1	1	
Steroids	0	0	0	0	1	1	0	1	
Alkaloids	1	1	1	1	1	1	1	1	
Coumarins	0	0	0	0	0	0	0	0	
Flavonoids	1	0	0	0	0	0	0	0	
Emodins	0	0	0	0	0	0	0	0	
Chloroform									
Tannins	1	1	1	1	1	1	1	1	
Saponins	1	1	1	1	1	1	0	0	
Terpenoids	1	1	1	1	1	1	1	1	
Steroids	0	0	0	0	0	0	1	0	
Alkaloids	0	0	0	0	0	0	0	0	
Coumarins	1	1	1	1	1	1	1	1	
Flavonoids	1	1	1	1	1	1	1	1	
Emodins	0	0	0	0	0	0	0	0	
Ethyl acetate									
Tannins	1	1	1	1	1	1	1	1	
Saponins	0	0	0	0	0	0	0	0	
Terpenoids	1	1	1	1	1	1	1	1	
Steroids	0	1	1	1	0	0	1	1	
Alkaloids	1	0	0	1	1	1	1	1	
Coumarins	1	1	1	1	1	1	1	1	
Flavonoids	1	1	1	1	1	1	1	1	
Emodins	0	0	0	0	1	1	0	0	
Sum of compound presences	12	11	11	12	14	15	13	14	

1 = presence; 0 = absence.

In the communities visited, infusions and preparations of these plants were used to treat stomach disorders, which could be related to the presence of these compounds. Similarly, Faizi and Naz [60] identified japhrin alkaloids with antimicrobial potential in *T. patula*.

CONCLUSION

Based on the exploration, documentation, collection and phytochemical evaluation of two species of *Tagetes* and two species of *Clinopodium*, it is concluded that *T. lucida*, *T. lunulata*, *C. macrostemum* and *C. mexicanum* are used for the treatment of disorders and gastrointestinal diseases, including ethno-diseases (e.g., 'empacho'), by indigenous communities of Oaxaca, Mexico. Among

species, significant differences were found in total polyphenol and flavonoid contents and antioxidant activity. The effect of species on the phenolic content and antioxidant activity was significantly higher than the effect of sample location or ecological-environment effect. *T. lucida* had the highest phenolic compound and antioxidant activity values, while *C. macrostemum* had the lowest. Tannins, alkaloids, steroids, terpenoids, and coumarins were detected in all the evaluated species, and emodins were detected only in *T. lucida*. One or more of these compounds have antioxidant and antimicrobial properties, thus validating the traditional use of these native species for treating disorders or stomach pain as indicated, in local terms, by the indigenous communities visited.

Conflicts of Interests

The authors have not declared any conflicts of interest.

ACKNOWLEDGMENTS

The authors confirm that all the communities mentioned in this study gave us their consent to perform the work and allowed us to conduct interviews and share parts of their knowledge.

REFERENCES

- Salmerón-Manzano E., Garrido-Cardenas J.A., Manzano-Agugliaro F. Worldwide research trends on medicinal plants. *Int J Environ Res Public Health*. 2020;17:10.
- Cai H., Zhang Y., Qiu Z., Luo P. Gene expression of antioxidant metabolic enzymes in grape extracts. *Cell Mol Biol*. 2021;67(6):200-212.
- World Health Organization (WHO). WHO global report on traditional and complementary medicine 2019. World Health Organization, Geneva. 2019.
- Touiti N., Houssaini T.S., Achour S. Overview on pharmacovigilance of nephrotoxic herbal medicines used worldwide. *Clinical Phytoscience*. 2021;7:1-9.
- Khan F., Pandey P., Singh A., Upadhyay T.K., AboElnaga S.M.H., Al-Majjar M.A.A., Saeed M., Kahrizi D. Unveiling antioxidant and antiproliferative effects of *Prosopis juliflora* leaves against human prostate cancer LNCaP cells. *Cell Mol Biol*. 2022;68(11):20-27.
- Zhang Q. Global situation and WHO strategy on traditional medicine. *Tradit Med Mod Med*. 2018;1:11-13.
- Lee E.L., Richards N, Harrison J, Barnes J. Prevalence of use of traditional, complementary and alternative medicine by the general population: A systematic review of national studies published from 2010 to 2019. *Drug Safety*. 2022;45:713-735.
- Sánchez-González A., Granados-Sánchez D., Simón-Nabor R. Uso medicinal de las plantas por los Otomíes del municipio de Nicolás Flores, Hidalgo, México. *Rev Chapingo Serie Horticult*. 2008;14:271-279.
- Gómez-Álvarez R. Plantas medicinales en una aldea del estado de Tabasco, México. *Rev Fitotec Mex*. 2012;35:43-49.
- Arrazola-Guendulay A.A., Hernández-Santiago E., Rodríguez-Ortiz G. Conocimiento tradicional de plantas silvestres en una comunidad de los Valles Centrales de Oaxaca. *Rev Mex Agroecosist*. 2018;5:55-78.
- Ortega-Cala L.L., Monroy-Ortiz C., Monroy-Martínez R., Colín-Bahena H., Flores-Franco G., Luna-Cavazos M., Monroy-Ortiz R. Plantas medicinales utilizadas para enfermedades del sistema digestivo en Tetela del Volcán, Estado de Morelos, México. *Bol Latinoam Caribe Plant Med Aromat*. 2019;18:106-129.
- García-Sánchez F., López-Villafranco M.E., Aguilar-Rodríguez S., Aguilar-Contreras A. Etnobotánica y morfo-anatomía comparada de tres especies de *Tagetes* que se utilizan en Nicolás Romero, Estado de México. *Bot Sci*. 2012;90:221-232.
- Mata-Pinzón S., Pérez-Ortega G., Reyes-Chilpa R. Plantas medicinales para el tratamiento del susto y mal de ojo. Análisis de sus posibles efectos sobre el Sistema Nervioso Central por vía transdérmica e inhalatoria. *Etnobiología* 2018;16:30-47.
- Pérez-Ortega G., González-Trujano M.E., Ángeles-López G.E., Brindis F., Vibrans H., Reyes-Chilpa R. *Tagetes lucida* Cav.: Ethnobotany, phytochemistry and pharmacology of its tranquilizing properties. *J Ethnopharmacol*. 2016;181:221-228.
- Domínguez-Vázquez G., Castro-Ramírez A.E. Usos medicinales de la familia Labiatae en Chiapas, México. *Etnobiología* 2002;2:19-31.
- Universidad Nacional Autónoma de México (UNAM). Biblioteca Digital de la Medicina Tradicional Mexicana. Universidad Nacional Autónoma de México (UNAM). 2009. Ciudad de México, México. <http://www.medicinatradicionalmexicana.unam.mx/>.
- Xu LW., Chen J., Qi HY., Shi Y.P. Phytochemicals and their biological activities of plants in *Tagetes* L. *Chin Herb Med*. 2012;4:103-117.
- Salehi B., Valussi M., Morais-Braga M.F.B., Pereira-Carneiro J.N., Borges L.A.L.A., Douglas CH.D., Vitalini S., Kręgiel D., Antolak H., Sharifi-Rad M., Cirone S.N.C., Yousaf Z., Martorell M., Iriti M., Carradori S., Sharifi-Rad J. *Tagetes* spp. essential oils and other extracts: chemical characterization and biological activity. *Molecules*. 2018;23:2847.
- Frezza C., Venditti A., Serafini M., Bianco A. Phytochemistry, Chemotaxonomy, Ethnopharmacology, and Nutraceuticals of Lamiaceae. In: Atta-ur-Rahman (ed.) *Studies in Natural Products Chemistry* Vol. 62, Elsevier, 2019, pp. 125-178.
- Aquino R., Cáceres A., Morelli S., Rastrelli L. An extract of *Tagetes lucida* and its phenolic constituents as antioxidants. *J Nat Prod*. 2002;65:1773-1776.
- Céspedes C.L., Avila J.G., Martínez A., Serrato B., Calderón-Mugica J.C., Salgado-Garciglia R. Antifungal and antibacterial activities of Mexican tarragon (*Tagetes lucida*). *J Agric Food Chem*. 2006;54:3521-3527.
- Wang W., Xu H., Chen H., Tai K., Liu F., Gao Y. *In vitro* antioxidant, anti-diabetic and antilipemic potentials of quercetagenin extracted from marigold (*Tagetes erecta* L.) inflorescence residues. *J Food Sci Technol*. 2016;53:2614-2624.
- González-Trujano M.E., Gutiérrez-Valentino C., Hernández-Aarámuro M.Y., Díaz-Reval M.I., Pellicer F. Identification of some bioactive metabolites and inhibitory receptors in the antinociceptive activity of *Tagetes lucida* Cav. *Life Sci*. 2019;231:116523.

24. Arroyo A.R., Chacón B., Maki K. Screening and selection of plants by positive pharmacologic effect on jejunum muscular contractility. *Pharm Biol.* 2004;42:24-29.
25. Pérez-Gutierrez R.M., Gallardo-Navarro Y.T. Antioxidant and hepatoprotective effects of the methanol extract of the leaves of *Satureja macrostema*. *Pharmacogn Mag.* 2010;6:125-132.
26. Villa-Ruano N., Zurita-Vásquez G.G., Pacheco-Hernández Y., Betancourt-Jiménez M.G., Cruz-Durán R., Duque-Bautista H. Anti-lipase and antioxidant properties of 30 medicinal plants used in Oaxaca, Mexico. *Biol Res.* 2013;46:53-160.
27. Alvarado A.M., García-Trejo F., Cardador-Martínez A., Magallán-Hernández F. *Clinopodium mexicanum*: potential and difficulties for the sustainable use of a Mexican medicinal plant. *Bol Latinoam Caribe Plant Med Aromat.* 2020;19:149-160.
28. Estrada-Reyes R., Martínez-Vázquez M., Gallegos-Solís A., Heinze G., Moreno J. Depressant effects of *Clinopodium mexicanum* Benth. Govaerts (Lamiaceae) on the central nervous system. *J Ethnopharmacol.* 2010;130:1-8.
29. Instituto Nacional de Estadística y Geografía (INEGI). (2008). Prontuario de información geográfica municipal de los Estados Unidos Mexicanos. Instituto Nacional de Estadística y Geografía, Mexico, 2008, https://www.inegi.org.mx/contenidos/app/mexicocifras/datos_geograficos/20/20239.pdf
30. IUSS Working Group WRB. World Reference Base for Soil Resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports No. 106. FAO, 2015.
31. Singleton V.L., Rossi J.A. Colorimetry of total phenolics with phosphomolybdic phosphotungstic acid reagents. *Am J Enol Vitic.* 1965;16:144-158.
32. Lin J.Y., Tang C.Y. Determination of total phenolic and flavonoid contents in selected fruits and vegetables, as well as their stimulatory effects on mouse splenocyte proliferation. *Food Chem.* 2007;101:140-147.
33. Zhishen J., Mengcheng T., Jianming W. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 1999;64:555-559.
34. Pękal A., Pyrzyńska K. Evaluation of aluminium complexation reaction for flavonoid content assay. *Food Anal Methods.* 2014;7:1776-1782.
35. Brand-Williams W., Cuvelier M.E., Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci Technol.* 1995;28:25-30.
36. Benzie I.F., Strain J.J. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol.* 1999;299:15-27.
37. Savithramma N., Linga-Rao M., Suhlulatha D. Screening of medicinal plants for secondary metabolites. *Middle-East J Scient Res.* 2011;8:579-584
38. Marcus A.C., Edori O.S., and Maduagu MC. Phytochemical and anti-microbial screening of *Phyllanthus fraternus* and *Taraxacum officinale* leaves. *Biochem Anal Biochem.* 2019;8:1-5.
39. SAS Institute Inc. (SAS). Base SAS ® 9.1.3 Procedures Guide, Second Edition, Volumes 1, 2, 3, and 4. Cary, NC: SAS Institute Inc., 2006.
40. Bello-González M.A., Salgado-Garciglia R. Plantas medicinales de la comunidad indígena Nuevo San Juan Parangaricutiro, Michoacán, México. *Polibotanica.* 2007;9:126-138.
41. Ortega-Ortega T., Vázquez-García V. *Satureja macrostema*: situación ambiental, conocimiento local y roles de género. *Madera y Bosques.* 2014;20:71-86.
42. Aguilar-Santelises R., del Castillo R.F. Demographic and socio-economic determinants of traditional plant knowledge among the Mixtecs of Oaxaca, Southern Mexico. *Hum Ecol.* 2015;43:655-667.
43. Rangel-Landa S., Casas A., Rivera-Lozoya E., Torres-García I., and Vallejo-Ramos M. Ixcatec ethnobotany: Plant management and biocultural heritage in Oaxaca, Mexico. *J Ethnobiol Ethnomed.* 2016;12:30.
44. Gómez-Luna R.E., Manzanero-Medina G.I., and Vázquez-Dávila M.A. Floristic and social aspects in zapotec orchards in Lachatao, Northern Sierra of Oaxaca, Mexico. *Rev Bio Ciencias.* 2017;4:1-15.
45. Tello-Ortega K.E., Hernández-Santiago E., Rodríguez-Ortiz G. Medicina alternativa complementaria en el tratamiento de enfermedades crónicas en el sur de Oaxaca, México. *Cienc Ergo sum.* 2020;27:1-10.
46. Cicevan R., Hassan M.A., Sestras A.F., Prohens J., Vicente O., Sestras R.E., Boscaiu. Screening for droughty tolerance in cultivars of the ornamental genus *Tagetes* (Asteraceae). *Peer J.* 2016;4:2133.
47. Chen J., Ning S., Lu X., Xiang W., Zhou X., Bu Y., Li L., Huang R. Variation in flavonoid and antioxidant activities of *Pyrrosia petiolosa* (Christ) Ching from different geographic origins. *Front Plant Sci.* 2023;14:1173489.
48. Zhang X.D., Yu Y.G., Yang D.F., Qi Z.C., Liu R.Z., Deng F.T., Caid Z.X., Li Y., Sun Y.F., Liang Z.S. Chemotaxonomic variation in secondary metabolites contents and their correlation between environmental factors in *Salvia miltiorrhiza* Bunge from natural habitat of China. *Ind Crops Prod.* 2018;113:335-47.
49. Ullah A., Munir S., Badshah S.L., Khan N., Ghani L., Poulson B.G., Emwas A-H., Jaremko M. Important flavonoids and their role as a therapeutic agent. *Molecules.* 2020;25:5243.
50. Núñez S., Moliner C., Valero M.S., Mustafa A.M., Maggi F., Gómez-Rincón C., López V. Antidiabetic and anti-obesity properties of a polyphenol-rich flower extract from *Tagetes erecta* L. and its effects on

- Caenorhabditis elegans* fat stoages. J Physiol Biochem. 2023;79:427-440.
51. Hernández T., Canales M., Flores C, García AM, Durán A, Avila J.G. Antimicrobial activity of *Tagetes lucida*. Pharm Biol. 2006;44:19-22.
52. Hernandez-Leon A., González-Trujano M.E., Narváez-González F., Pérez-Ortega G., Rivero-Cruz F., Aguilar M.I. Role of β -caryophyllene in the atinociceptive and anti-inflammatory effects of *Tagetes lucida* Cav. essential oil. Molecules. 2020;25:1-15.
53. Li X., Li L., Lei W., Chua H.Z., Li Z., Huang X., Wang Q., Li N., Zhang H. Traditional Chinese medicine as a therapeutic option for cardiac fibrosis: pharmacology and mechanisms. Biomed Pharmacother. 2021;142:111979.
54. Morales-Torres K., Mora-Román J.J., Fallas-Ramírez J.M., Rodríguez-Arrieta J.A., Orozco-Aguilar J. Absence of nephroprotective effect: evaluation of an aqueous extract of *Clinopodium vimineum* (L.) Kuntzie (Lamiaceae) a renal failure model with female Wistar rats. Eur J Pharm Med Res. 2022;9(10):43-50.
55. Dunkić V., Kremer D., Grubescic R.j., Rodriguez J.V., Ballian D., Bogunic F., Stesevic D., Kosalec I., Bezic N., Stabentheiner E. Micromorphological and phytochemical traits of four *Clinopodium* L. species (Lamiaceae). S Afr J Bot. 2017;111:232-241.
56. Li W., Gao Y., Zhao J., Wang Q. Phenolic, flavonoid, and lutein ester content and antioxidant activity of 11 cultivars of Chinese. J Agric Food Chem. 2007;55:8478-8484.
57. Monterrosas-Brisson N., Herrera-Ruiz M., Jiménez-Ferrer E., Bahena-Pérez R., Avilés-Flores M., Fuentes-Mata M., Martínez-Duncker I., González-Cortazar M. Anti-inflammatory activity of coumarins isolated from *Tagetes lucida* Cav. Nat Prod Res. 2020;34:3244-3248.
58. Estrada-Soto S., González-Trujano M.E., Rendón-Vallejo P., Arias-Durán L., Ávila-Villarreal G., Villalobos-Molina R. Antihypertensive and vasorelaxant mode of action of the ethanol-soluble extract from *Tagetes lucida* Cav. aerial parts and its main bioactive metabolites. J Ethnopharmacol. 2021;266:113399.
59. Villa-Ruano N., Pacheco-Hernández Y., Cruz-Durán R., Lozoya-Gloria E. Volatiles and seasonal variation of the essential oil composition from the leaves of *Clinopodium macrostemum* var. *laevigatum* and its biological activities. Ind Crop Prod. 2015;77:741-747.
60. Faizi S., Naz A. Jafrine, a novel and labile β -carboline alkaloid from the flowers of *Tagetes patula*. Tetrahedron. 2002;58:6185-6197.