

Foliar and pollen grain micromorphology of some species of *Astragalus*

Received: 06.10.2023 ===== Revised: 13.11.2023 ===== Accepted: 15.11.2023

Sedigheh Eskandari: MSc Graduate, Department of Plant Biology, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran**Azadeh Akhavan Roofigar:** Research Assistant Prof., Department of Natural Resources, Isfahan Agricultural and Natural Resources Research Center, Agricultural Research, Education and Extension Organization (AREEO), Isfahan, Iran**Shahrokh Kazempour-Osaloo**✉: Prof., Department of Plant Biology, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran (skosaloo@modares.ac.ir)**Abstract**

The present study is focused on identifying the differences between the closely related sections *Microphysa* and *Campylanthus* in the genus *Astragalus* using micromorphological characteristics. For this purpose, the morphological characteristics of leaf epidermal cells and stomata as well as pollen grains have been studied. The results obtained from light- and scanning electron microscopy (LM and SEM) indicated that, stomatal type was typically anemocytic in most species, while the anisocytic type was only found in two species, viz. *A. callistachys* and *A. microphysa*. The epidermal cells in most species were observed to have a polygonal shape with straight anticlinal walls. However, in *A. cephalanthus*, both polygonal and irregular cell shapes as well as sinuate anticlinal walls were observed. All pollen grains shed as monad and exhibited small size, single shape, tricolporate aperture type, isopolarity and symmetry, with rounded or elongated pores. It was concluded that, the micromorphological data alone were not enough to separate these two sections and determine the limits of the species.

Keywords: *Campylanthus*, epidermal cells, *Fabaceae*, *Microphysa*, pollen grain**ریزریخت‌شناسی برگ و دانه گرده برخی گونه‌های *Astragalus****

دریافت: ۱۴۰۲/۰۷/۱۴ ===== بازنگری: ۱۴۰۲/۰۸/۲۲ ===== پذیرش: ۱۴۰۲/۰۸/۲۴

صدیقه اسکندری: دانش‌آموخته کارشناسی ارشد گروه علوم گیاهی، دانشکده علوم زیستی، دانشگاه تربیت مدرس، تهران، ایران
آزاده اخوان روفیگر: استادیار پژوهش بخش تحقیقات منابع طبیعی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی استان اصفهان، سازمان تحقیقات، آموزش و ترویج کشاورزی، اصفهان، ایران

شاهرخ کاظم‌پور اصلو✉: استاد گروه علوم گیاهی، دانشکده علوم زیستی، دانشگاه تربیت مدرس، تهران، ایران (skosaloo@modares.ac.ir)

خلاصه

مطالعه حاضر بر شناسایی تفاوت‌های بین بخش‌های به هم نزدیک *Microphysa* و *Campylanthus* در جنس *Astragalus* (باقلابیان) با استفاده از ویژگی‌های ریزریخت‌شناسی متمرکز است. به این منظور، ریخت‌شناسی سلول‌های اپیدرمی و روزنه‌های برگ و دانه‌های گرده در هفت گونه بررسی شد. نتایج حاصل از مطالعات میکروسکوپی (نوری و الکترونی) نشان داد که نوع روزنه در بیشتر گونه‌ها به طور مشخص آنموسیتیک بود، در حالی که روزنه آنیزوسیتیک تنها در دو گونه *A. microphysa* و *A. callistachys* یافت شد. سلول‌های اپیدرمی در بیشتر گونه‌ها به شکل چندضلعی منظم با دیواره صاف مشاهده شدند، اما در گونه *A. cephalanthus* سلول‌های اپیدرمی به هر دو شکل چندضلعی و نامنظم و همچنین دیواره عمودی به شکل سینوسی دیده شد. براساس یافته‌های این مطالعه، پراکنش تمام دانه‌های گرده به طور منفرد، دارای اندازه کوچک و سه شیار منفذدار، جورقطب و متقارن با منافذ گرد یا کشیده بودند. طبق نتایج به دست آمده، داده‌های ریزریخت‌شناسی به تنهایی برای جداسازی این دو بخش و تعیین حدود گونه‌ها، معیار کافی تشخیص داده نشدند.

واژه‌های کلیدی: باقلا بیان، دانه گرده، سلول‌های اپیدرمی، *Microphysa*، *Campylanthus*

* مستخرج از پایان‌نامه کارشناسی ارشد نگارنده نخست به راهنمایی دکتر شاهرخ کاظم‌پور اصلو و دکتر آزاده اخوان روفیگر ارائه شده به دانشگاه تربیت مدرس

Introduction

Astragalus L. (*Fabaceae*) is the largest genus of flowering plants with approximately 3156 species classified into 255 sections (Maassoumi 2020). *Astragalus* is known for its adaptation to various ecological conditions and habitat. The plant can be found in cool to warm climates, semi-arid to arid regions, and mountainous areas, which has resulted in a wide range of forms such as annual, perennial, herbaceous thorn cushions form (Castillon *et al.* 2020). The genus *Astragalus* displays a wide range of morphological diversity, such as differences in vegetative form and habitat, plant size, types of hairs (Ghahremaninejad 2004), inflorescence type, length of inflorescence peduncle and raceme, petal length, and other characteristics (Podlech & Zarre 2013). The diversity of ecological factors and changes in habitat can also result in variation among *Astragalus* species. These variations include preferences for different growing environments such as mountain slopes, flatlands, riverbanks, agricultural fields with dry soil, and barren lands with hard, stony or sandy soils. *Astragalus* species exhibit a high level of sensitivity to regional and microclimatic conditions (Maassoumi 2020). Although *Astragalus* has a long biogeographical history, our understanding of its regional connections and distribution in the Old World is still limited. Until now, there is no comprehensive, up-to-date, and integrated information available on diverse spatial distribution patterns in a broad continental-scale framework (Zarre & Azani 2013).

With more than 848 species in 70 sections, Iran is a major center of diversity for the *Astragalus* genus, representing more than 10% of the country's plant species. (Maassoumi 2005, Bagheri *et al.* 2014, Ghahremaninejad *et al.* 2022). Of these species, over 570 are endemic to the flora of Iran (Maassoumi 2005). The presence of endemic species with a wide distribution suggests that, their center of origin is mainly in the Zagros Mountains and to a lesser extent in the Alborz Mountains or scattered throughout other high-altitude areas in the central plateau (Ghahremaninejad *et al.* 2016,

Maassoumi 2005, 2020, Noroozi *et al.* 2020, Mahmoodi *et al.* 2022).

Modern taxonomists often use palynology as a method to distinguish closely related species (Khan *et al.* 2018, Ullah *et al.* 2018 Amina *et al.* 2020). For micromorphological studies, the Scanning Electron Microscope (SEM) is an advanced approach that enables the examination of pollen grain structures in exceptional detail (Bagheri *et al.* 2019, Majeed *et al.* 2020, Umber *et al.* 2022). SEM has been employed in many plant families, including *Fabaceae* (Ekici *et al.* 2005, Lashin, 2006, Büyükkartal *et al.* 2012, Bagheri *et al.* 2019), *Asteraceae* (Karbalaie *et al.* 2021, Khan *et al.* 2021, Nabila *et al.* 2022, Umbe *et al.* 2022), *Caryophyllaceae* (Ullah *et al.* 2019), *Acanthaceae* (Raza *et al.* 2020), *Cactaceae* (Majeed *et al.* 2020), and *Lamiaceae* (Bahadur *et al.* 2022) due to its high resolution and ability to accurately display the surface of pollen grains.

Despite numerous studies that have attempted to classify the diverse and polymorphic genus *Astragalus*, some classification issues remain unresolved (Khan *et al.* 2022). To address this, micromorphological characteristics of pollen grains have been introduced as a valuable tool in the taxonomic classification of the genus (Yasmin *et al.* 2010, Khan *et al.* 2022). However, only few sections of *Astragalus* species have been studied using pollen analysis, including sect. *Hololeuce* Bunge (Ceter *et al.* 2013, Uzun *et al.* 2021), sect. *Onobrychoidei* DC. (Pinar *et al.* 2009), sect. *Malacothrix* Bunge (Oskouian *et al.* 2007), sect. *Hymenostegis* Bunge (Bagheri *et al.* 2019) and sect. *Macrophyllum* Boiss. (Ranjbar *et al.* 2012). The use of pollen grains in systematics has proven to be valuable due to the information that can be obtained from their characteristics such as their shape, size, pores, modes of attachment, and symmetry. These characteristics aid in distinguishing and classifying closely related species within a genus. *Papilionoideae*, a subfamily of the *Fabaceae*, displays a great degree of variability in the size, arrangement of pores, and surface decorations of its pollen grains, which is referred to as Eurypalynous. The shape of pollen grains is usually radially symmetrical and free, with a

tricolporate arrangement that is occasionally colpate or porate. The shape of the grains is typically prolate or almost subprolate, and the tectum is mostly reticulate, with some species displaying fossulate, foveolate, or fossulate-rugulate tectum. However, more than 70% of the studied taxa exhibit tricolporate pollen grains with reticulate tectum (Perveen & Qaiser 1998). Changes in the exine decorations, grooves, and pattern of pollen grains are significant and contribute to their differentiation and classification in the subfamily.

The micromorphological diversity of epidermal cells is one of the main sources of data in taxonomy and plays a fundamental role in systematics (Metcalf 1988). The foliar epidermal cells of *Astragalus* are remarkable characters for the distinction of different species (Hayat *et al.* 2009). These characters include the size and shape of epidermal cells, distribution of stomata, size of guard cells, and number of subsidiary cells. The use of foliar anatomical attributes in plant systematics has been over the last 100 years (Bahadur *et al.* 2019). Therefore, due to the widespread distribution of this huge genus and for its complex taxonomy, the micromorphological characteristics of foliar epidermal in some selected species of *Astragalus* were monitored with SEM techniques to better understanding the taxonomic value of micromorphological traits among different species of *Astragalus* and determine the influence these traits to resolving taxonomic conflicts of this genus.

This study focuses on *Astragalus* sect. *Microphysa* and sect. *Campylanthus*. Different analyses of molecular phylogenetic data have shown that, these sections are not monophyletic; instead, their species are intermixed. In addition, these two sections are identified as sister taxa and are nested within a polytomy clade (Kazempour Osaloo *et al.* 2003, 2005, Bagheri *et al.* 2017). In this study, seven species of *Astragalus* from *Microphysa* and *Campylanthus* sections were investigated in terms of micromorphology of foliar epidermal cells and pollen grain. The aim of this study was to examine the micromorphology of reproductive and vegetative parts in seven species of *Astragalus* belonging to the *Microphysa* and *Campylanthus* sections. The study

aimed to investigate whether the micromorphological characteristics could be used to distinguish between different species and sections.

Materials and Methods

- Sampling

All leaf and pollen samples were obtained from 24 specimens belonging to seven species (at least three specimens per species were studied) from *Microphysa* and *Campylanthus* sections deposited at the herbarium of the Isfahan Agricultural and Natural Resources Research Center (SFAHAN), Isfahan (Iran), listed in table 1.

- Light microscopy

The dried samples were fixed in a solution of ethanol and acetic acid (70:30) for 24–48 hr. They were then washed with distilled water and immersed in a solution of hydrogen peroxide and acetic acid (1:1) for 40–45 min at 50–60 °C. The samples were re-washed with distilled water and stained with Carmine for 2–3 min if necessary. The epidermis was separated and examined under a Light Microscope (LM), and various photographs were taken using slides.

- Scanning electron microscopy

Scanning Electron Microscope (SEM) is widely used to observe surface structures. Micromorphological studies were conducted using the SEM device to investigate the upper epidermis of the leaf and pollen grains (Table 1) from the studied species. The specifications of the SEM device (AIS2300C SERON) used in the laboratory of Isfahan University of Technology, Faculty of Materials (Isfahan, Iran) were with a maximum length and width of 2.5 cm, and a maximum height of 3 cm. To prepare the sample, 10 nm layer of gold was coated on the surface of the samples, and then transferred to the SEM device for imaging. After many samples of leaf epidermis and stomata were photographed to record the results, the best pictures were selected and the parameters were measured and recorded using Digimizer 5.4.9 software.

Table 1. Information of examined *Astragalus* sect. *Microphysa* and sect. *Campylanthus* in Isfahan Province

No.	Taxon	Locality	Altitude (m)	Collector	Voucher No.
1	<i>A. argyrostachys</i>	Semirom, Garmook	2350	Feyzi	13543■
		Fereydan, Aghche	2300	Feyzi	12267■◆◆
		Isfahan prov.: Tiran to Damaneh, Tang-e-kolang	2500	Feyzi & Shams	12514■◆◆
2	<i>A. callistachys</i>	20 km Shahreza to Semirom	2100	Nowroozi & Etemadi	526■◆◆
		Tiran, Ghameshloo	2100	Nowroozi & Feyzi	5336■◆
		5 km Delijan to Tehran	1450	Nowroozi	4618■◆
		Najaf-abad, Jannat-abad	1600	Nowroozi	492■
3	<i>A. campylanthus</i>	Semirom, Vanak, Dalan-kooch	2200	Nowroozi & Bozorgi	3597■◆◆
		Meymeh	2400	Feyzi	13193■
		Dehaghan, Kheir-abad	2200	Feyzi	6755■●
		Fereydan, Aghcheh	2300	Feyzi	12269■◆
4	<i>A. cephalanthus</i>	Semirom, Hanna	2250	Nowroozi	4005■◆◆
		Chal-e-siah to Hosein-abab	1850	Nowroozi & Feyzi	6056■◆
		Meymeh	2400	Nowroozi	13191■
		Boroोजen to Mobarakeh	2100	Nowroozi & Feyzi	8026■◆
		Semirom to Padena	2600	Mohajeri	7887■
		Chadegan to Bidak	2500	Nowroozi & Etemadi	1059■
		Fereydan	2400	Feyzi & Shams	12466■
		Chadegan, Zayandeh-rood	2200	Nowroozi & Matin	6147●
5	<i>A. microphysa</i>	Chal-e-siah	1850	Nowroozi	6055■◆
		Kashan to Ghamsar	1250	Nowroozi	13565■◆◆
		Buein, Tang-e-doozan	2800	Nowroozi & Etemadi	1305■◆
		Meymeh to Muteh	1920	Feyzi	13108◆
		Najaf-abad, Dehagh	2100	Feyzi & Shams	12582●
		Zarrin-shahr	1750	Feyzi	14644●
6	<i>A. susianus</i>	Fereydoon-shahr, Sang-baran	2400	Feyzi & Asfa	10404■
		Khansar	2700	Feyzi	8539■
		Buein-ghaleh	2550	Feyzi & Saeidfar	6045■◆
		Semirom, Hanna	2400	Feyzi & Shams	14370◆
		Daran	2300	Feyzi & Saeidfar	6415●
		Fereydan, Dalan-kooch	2400	Feyzi & Eftekhari	11158●
7	<i>A. ruterianus</i>	Fereydan, Ghal-e-Shahrokh	2500	Nowroozi & Yazdani	13323◆◆

■: Specimens used for LM, ◆: Specimens used for SEM from epidermis of the leaf, ●: Specimens used for SEM from pollen grains

To prepare the pollen sample, the anthers are first separated from the flower. The terminal anther is then slowly crushed on a plate with a needle to create a powder. Some of the powdered anthers are carefully and delicately placed on the electron microscope plates and observed after completing the necessary preparation steps. After taking multiple pictures of pollen grain samples with SEM, the best pictures were selected, and the parameters were measured and recorded using Digimizer 5.4.9 software (Table 5). The micro-morphological analysis involved the measurement of 26 characters for each species. Principal Coordinate Analysis (PCoA) was conducted using GenAlex.

Results

- Epidermal cells and stomata

Different foliar epidermal features and the results of qualitative and quantitative analysis of the studied species under LM and SEM are summarized in tables 2–4. Table 2 pertains to LM images, while tables 3 and 4 pertain to SEM images. The shape of epidermal cells was more specifically investigated, and the type of stomata was determined using LM, while SEM was more suitable for determining quantitative traits, such as measuring the length and width of epidermal cells and stomata. The results obtained from LM and SEM indicates that, stomatal type is typically anemocytic in

most species, while the anisocytic type is found only in two species, viz. *A. callistachys* and *A. microphysa*.

The shape of epidermal cells can be divided into two main categories: regular (polygonal) and irregular. In addition, based on the shape of the anticlinal wall, the cells of the epidermis can be classified into three forms: sinuate, undulate, and straight. In the images obtained from both light and electron microscopes, the epidermal cells in most species were observed to have a polygonal shape with straight anticlinal walls. However, in *A. cephalanthus*, both polygonal and irregular cell shapes were

observed, and sinuate anticlinal walls were also present in this species. Based on these observations, the studied species can be divided into three general groups based on the shape of the anticlinal wall and epidermal cells: 1. Polygonal cells with straight walls: *A. argyrostachys* (Figs 1A, 2A), *A. callistachys* (Figs 1D, 2B), *A. campylanthus* (Figs 1B, 2C), and *A. susianus* (Figs 1C, 3C), 2. Polygonal cells with undulate walls: *A. reurianus*: (Fig. 3B), *A. microphysa* (Figs 1F, 3A), and *A. cephalanthus* (Figs 1E, 2D), and 3. Irregular cells with sinuolate walls: *A. cephalanthus* (Figs 1E, 2D).

Table 2. Measurement of quantitative traits of epidermis and stomata with LM

No.	Taxon	Herbarium No.	Average epidermal cell (μm)		Average stomata (μm)		Average stomatal guard cell (μm)		Average stomatal aperture (μm)	
			Length	Width	Length	Width	Length	Width	Length	Width
1	<i>A. argyrostachys</i>	13543	43	23.6	22	21	20	6.8	11.9	3.2
2		12267	40	18.6	22	21	20.8	7.6	11	4.7
3		12514	39.2	18	21.9	17.2	20.3	4.6	12	4
4	<i>A. callistachys</i>	526	34.1	19.5	20	20.5	17.5	6.9	8	4.5
5		5336	43	25.5	24.2	25.3	24.7	9.5	11	5
6		4618	38	20	24	21	20.5	7.2	10.5	4.2
7		492	39	18.7	19.2	17.5	16.5	6.8	10	4.1
8	<i>A. campylanthus</i>	3597	32.9	18	17.2	17	17.6	4.4	11.3	3.9
9		13193	34.9	20.1	23	20.6	19.4	4.6	9.2	3.4
10		6755	35.4	17.7	22.7	18.8	20	7.8	9.7	3.5
11		12269	32.8	18.4	18.4	18.8	17.5	7.9	8.9	3.7
12		<i>A. cephalanthus</i>	4005	54.6	27.8	24	17	21	6	12.2
13	6056		50.4	22.1	24.3	19	20	5.4	9.8	5
14	13191		51.4	23.7	22.4	19	22	7.5	11	5
15	8026		55	28	25.4	18	21.7	7.8	12.8	4.8
16	7887		49	21.8	24	17	21.6	7	12	4.9
17	1059		58	34.6	22.5	18	21.2	7.7	11	3.8
18	12466		47	23.8	18	15.5	18	6.5	10.5	3.8
19	<i>A. microphysa</i>	12582	45	12	13	12	14.5	7	12.5	2.2
20		6055	25	9.7	10	8.4	10	3.2	6	2.1
21		14219	42	16	23	22	20	7.5	8	4.3
22		13565	34	20	20	13	18.2	5.4	8.8	3.7
23	<i>A. susianus</i>	10404	40	14	22	21	18	5.8	12	5
24		6405	37	13	21.7	18	17	4.2	10.5	3.5
25		8539	37	15.3	20	18	18	4.7	10	3.5

Table 3. Quantitative and qualitative characteristics of epidermis with SEM


No.	Taxon	Herbarium No.	Mean quantitative characteristics of epidermis		Qualitative characteristics of epidermis	
			Epidermal cell length (μm)	Epidermal cell width (μm)	Epidermal cell shape	Margin
1	<i>A. argyrostachys</i>	12267	54	22	Polygonal	Straight
2		12514	49	22.8	Polygonal	Straight
3	<i>A. callistachys</i>	526	35.5	17.1	Polygonal	Straight
4		4618	35.8	15.3	Polygonal	Straight
5		5336	39.3	20.3	Polygonal	Straight
6	<i>A. campylanthus</i>	3597	24.7	9.5	Polygonal	Straight
7		12269	28.7	10.8	Polygonal	Straight
8	<i>A. cephalanthus</i>	4005	45.4	22.5	Irregular	Sinuolate
9		6056	35	8.5	Polygonal	Undulate
10		8026	38.4	14.1	Polygonal	Undulate
11	<i>A. microphysa</i>	1305	37.6	13.9	Polygonal	Straight
12		6055	40	16	Polygonal	Straight
13		13108	50	25.5	Polygonal	Straight
14		13565	42.5	20.6	Polygonal	Straight
15	<i>A. ruterianus</i>	13323	40.7	10.5	Polygonal	Undulate
16		14370	30.5	9.5	Polygonal	Undulate
17	<i>A. susianus</i>	6405	38.2	9.2	Polygonal	Straight

Table 4. Quantitative and qualitative traits of stomata with SEM

No.	Taxon	Herbarium No.	Mean quantitative traits of stomata							Qualitative traits of stomata
			Stomata length (µm)	Stomata width (µm)	Ambiance of stomata	Stomatal guard cell length (µm)	Stomatal guard cell width (µm)	Stomatal pore length (µm)	Stomatal pore width (µm)	Type of stomata
1	<i>A. argyrostachys</i>	12267	22.1	17.8	60.6	23	5.4	12	5.6	Anemocytic
2		12514	20.1	13.78	52.5	20.2	4.8	11.5	5	Anemocytic
3	<i>A. callistachys</i>	526	22.8	13	56.3	23	4	9.5	4.5	Anisocytic
4		4618	18.3	10.5	46	18.8	3.9	10.7	4.4	Anisocytic
5		5336	23.5	16	61	15	2.9	11.3	5.3	Anisocytic
6	<i>A. campylanthus</i>	3597	20	17.4	51	17.5	3.5	10.4	4.1	Anemocytic
7		12269	17.1	11.5	47.1	14	3.5	9.4	4.4	Anemocytic
8	<i>A. cephalanthus</i>	4005	25.8	13	61.8	19.3	4.7	12.5	5.1	Anemocytic
9		6056	23.1	15	60.6	20	4.7	13.5	5.5	Anemocytic
10		8026	25.8	15.9	65	23	5.1	12.3	5	Anemocytic
11	<i>A. microphysa</i>	1305	20.1	15.2	56.3	16.5	4.1	7.4	3.5	Anemocytic
12		6055	18.5	13.2	49.2	15	4	9	3.1	Anisocytic
13		13108	24.7	11.2	59	19.5	3.8	10.3	5.3	Anemocytic
14		13565	20.4	13.1	52.9	19	5	9.1	3.2	Anemocytic
15	<i>A. ruterianus</i>	13323	15.7	7.5	37.9	14.5	2.8	7.5	3.1	Anemocytic
16		14370	16	8.4	39.3	15	2.5	9.3	3	Anemocytic
17	<i>A. susianus</i>	6405	21	17	64	16	3.8	10.7	4.2	Anemocytic

Table 5. Data related to the measured characteristics of pollen grains in the studied species

No.	Taxon	Herbarium No.	Mean quantitative trait			Mean qualitative trait		
			Polar axis (P) (μm)	Equatorial axis (E) (μm)	(P/E)	Shape	Type and number of pore/furrow	Type of decoration
1	<i>A. argyrostachys</i>	12267	29	25	1.16	Subprolate	Tricolporate	Rugulate
2		12514	22.5	12	1.87			
3	<i>A. callistachys</i>	526	27.5	13.2	2.08	Prolate	Tricolporate	Rugulate
4	<i>A. campylanthus</i>	3597	27	13	2.07	Prolate	Tricolporate	Rugulate
5		6755	25.1	11.1	2.26			
6	<i>A. cephalanthus</i>	6147	21.6	21.2	1.01	Spheroidal to prolate	Tricolporate	Reticulate
7		4005	26.3	13.4	1.96			Rugulate
8	<i>A. microphysa</i>	12582	28.5	24.5	1.16	Subprolate	Tricolporate	Reticulate
9		13565	33	23.5	1.4			
10		13644	26.1	23.2	1.12			
11	<i>A. ruterianus</i>	13323	19.5	16.1	1.21	Subprolate	Tricolporate	Reticulate
12	<i>A. susianus</i>	6415	24	20.1	1.19	Subprolate	Tricolporate	Reticulate
13		11158	23.4	20.2	1.16			

**Fig. 1.** LM of foliar epidermal cells and stomata of different *Astragalus* species: A. *A. argyrostachys*, B. *A. campylanthus*, C. *A. susianus*, D. *A. callistachys*, E. *A. cephalanthus*, F. *A. microphysa*.




Fig. 2. SEM of foliar epidermal cells and stomata of different *Astragalus* species: A1–A3. *A. argyrostachyes*, B1–B3. *A. callistachys*, C1–C3. *A. campylanthus*, D1–D3. *A. cephalanthus*.

The electron microscope measurements showed that, the longest stomatal length was found in the *A. cephalanthus* with an average of 25 μm , while the shortest was observed in the *A. callistachys* with an average of 16 μm . Based on the measurements from electron microscope images, it was found that, the *A. cephalanthus* has the widest stomatal aperture with an average of 14.1 μm , while the *A. callistachys* has the narrowest stomatal aperture with an average of 8–10 μm . The longest stomatal guard cell length was observed in the *A. cephalanthus* with an average of 21 μm , while the shortest length was found

in *A. microphysa* and *A. campylanthus* with approximately 16 μm . Among the examined taxa, *A. cephalanthus* had the widest stomatal guard cell, measuring 13 μm , while *A. campylanthus*, and *A. callistachys* had the narrowest, measuring around 3.5 μm . The stomatal pores of *A. cephalanthus* were also the longest, with a mean length of 13 μm , whereas those of *A. microphysa* were the shortest, with a mean length of 9 μm . When it comes to stomatal pore width, *A. cephalanthus* had the largest, measuring 3.5 μm , whereas *A. microphysa* had the smallest, also measuring 3.5 μm .




Fig. 3. SEM of foliar epidermal cells and stomata of different *Astragalus* species: A1–A3. *A. microphysa*, B1–B3. *A. ruterianus*, C1–C3. *A. susianus*.

- Pollen grains scanning electron microscopy

Pollen grains shed from the anthers were examined using SEM. Ideally, the pollen grains that were individually placed were photographed with higher magnification. Based on observations, all pollen grains exhibited small size, single shape, tricolporate aperture type, isopolarity and symmetry,

with rounded or elongated pores (Figs 4–5). Based on the calculations, the ratio of the polar axis (P) to the equatorial axis (E) was greater than one and ranged from 1.01 in *A. cephalanthus* from the sect. *Campylanthus* to 2.26 in the *A. campylanthus* from the sect. *Microphysa*.




Fig. 4. SEM of pollen grain of different *Astragalus* species: A1–A2. *A. argyrostachyes*, B1–B2. *A. callistachys*, C1–C2. *A. campylanthus*.




Fig. 5. SEM of pollen grain of different *Astragalus* species: A1–A2. *A. cephalanthus*, B1–B2. *A. microphysa*, C1–C2. *A. ruterianus*, D1–D2. *A. susianus*.

Discussion

Upon investigation of the quantitatively defined traits, no significant difference was observed between the species of *Microphysa* and *Campylanthus* sections. However, among the species studied, *A. cephalanthus* was found to be significantly larger than other species in terms of stomatal length and width, stomatal pore, and guard cells. The minimum and maximum length and width of epidermal cells were found in *A. campylanthus* and *A. agrostachys*, respectively, both of which belong to sect. *Campylanthus*. Therefore, it can be concluded that, the size of epidermal cells is highly variable among different species and cannot be used to differentiate between the species of the two mentioned sections. Based on observations, stomata in the sect. *Microphysa* were predominantly anemocytic, with some instances of anisocytic types, while all stomata in the sect. *Campylanthus* were anemocytic.

In terms of the anticlinal wall shape, species in the sect. *Microphysa* exhibited straight, undulate, and occasionally sinuolate shapes, while species in the sect. *Campylanthus* exhibited straight and occasionally undulate shapes. The type of pollen grain surface decoration observed in the samples can be broadly categorized as rugulate, mostly observed in species belonging to the sect. *Campylanthus*, and reticulate, observed in species of the sect. *Microphysa*. This suggests that pollen grain surface decoration can be an effective taxonomic trait for differentiating between the species of the two sections. Pollen grain shape was found to be broadly

categorized into three groups: almost spheroidal in *A. cephalanthus*, subprolate in *A. ruterianus*, *A. susianus*, *A. agrostachys* and *A. microphysa*, and prolate in the *A. callistachys* and *A. campylanthus*. In general, species belonging to the sect. *Campylanthus* exhibited subprolate and prolate shapes, while those in the sect. *Microphysa* exhibited prolate, subprolate, and spheroidal shapes. This is in congruence to other pollen studies in *Astragalus* species, which found the pollen in their study groups to be mainly prolate and subprolate (Pinar *et al.* 2009, Osman *et al.* 2014).

As mentioned earlier, the shape of the pollen grain is more or less variable, especially in the sect. *Microphysa*. Therefore, it cannot be considered as a significant characteristic in separating the species of the two sections. Other characteristics, such as stomatal features and epidermal cell size, seem to be more effective in distinguishing between the species of these sections. These results are also depicted in figure 6 as part of the PCoA analysis. It should be mentioned that, the results obtained from the set of micromorphological characters of pollen and epidermis are not consistent with the morphological characteristics of the two studied sections. The findings indicate that these characters were not efficient enough to distinguish between the two sections and their respective species. This suggests that, these characteristics may not be particularly useful in distinguishing between different species within the genus *Astragalus*. Other morphological or genetic traits may need to be considered in order to accurately differentiate between closely related species within the genus.




Fig. 6. Principal coordinate analysis (PCoA) score plot expressing the micromorphological variation (epidermis & pollen) of seven species of *Astragalus* sect. *Microphysa* and sect. *Campylanthus*.

Conclusion

The micromorphological traits of pollen grain, epidermal cells and stomata were studied to distinguish between the closely related *Microphysa* and *Campylanthus* sections in the genus *Astragalus*. The results showed that, these traits were not efficient enough to separate the two sections and their species. Therefore, it is necessary to use a combination of micromorphological and molecular approaches for accurate classification of the genus *Astragalus*. Micromorphological studies can save time and budget in the first

step, but genetic studies are required for validation and completion of the results.

Acknowledgments

The authors would like to express their gratitude to the Tarbiat Modares University (Tehran, Iran) authorities for approving and financing this project, and the assistance of the Agricultural and Natural Resources Research Center of Isfahan (Isfahan, Iran) for the permission to use the herbarium specimens. The authors also thank Ms. Karbalaei for her technical assistance.

References

- Bagheri, A., Akhavan Roofgar, A., Abbasi, S., Maassoumi, A.A., Rutten, T. & Blattner, F.R. 2019. Pollen morphology of *Astragalus* section *Hymenostegis* (Fabaceae) and evaluation of its systematic implications. *Grana* 58(5): 328–336.
- Bagheri, A., Maassoumi, A.A., Rahiminejad, M.R., Brassac, J. & Blattner, F.R. 2017. Molecular phylogeny and divergence times of *Astragalus* section *Hymenostegis*: An analysis of a rapidly diversifying species group in Fabaceae. *Scientific Reports* 7(1): 14033.
- Bagheri, A., Rahiminejad, M.R. & Maassoumi, A.A. 2014. A new species of the genus *Astragalus* (Leguminosae-Papilionoideae) from Iran. *Phytotaxa* 178(1): 38–42.
- Bahadur, S., Taj, S., Ahmad, M., Zafar, M., Gul, S., Shuaib, M., Butt, M.A., Hanif, U., Nizamani, M.M., Hussain, F. & Romman, M. 2022. Authentication of the therapeutic Lamiaceae taxa by using pollen traits observed under scanning electron microscopy. *Microscopy Research and Technique* 85(6): 2026–2044.
- Buyukkartal, H.N., Çolgeçen, H., Pinar, N.M. & Erdoğan, N. 2012. Seed coat ultrastructure of hard-seeded and soft-seeded varieties of *Vicia sativa*. *Turkish Journal of Botany* 37(2): 270–275.
- Castillon, E.E., Quintallina, J.A.V. & Salinas, A.D. 2020. Morphological range changes in Mexican *Astragalus radicans* Hornem. (Fabaceae: Galegeae): review of its taxonomy and nomenclature. *Phytotaxa* 470(2): 123–132.

- Ceter, T., Ekici, M., Pinar, N.M. & Ozbek, F. 2013. Pollen morphology of *Astragalus* L. section *Hololeuce* Bunge (Fabaceae) in Turkey. *Acta Botanica Gallica* 160(1): 43–52.
- Ekici, M., Yuzbas, D. & Aytac, Z. 2005. Morphology, pollen, seed structure and karyological study on *Astragalus ovalis* Boiss. and Balansa (sect. *Ammodendron*) in Turkey. *International Journal of Botany* 1(1): 74–78.
- Ferguson, I.K. & Skvarla, J. 1981. The pollen morphology of the subfamily Papilionoideae (Leguminosae). Pp. 859–896. *In*: Polhill, R.M. & Raven, P.H. (eds), *Advances in Legume Systematics*. Royal Botanic Garden, Kew.
- Ghahremaninejad, F. 2004. Value of trichome characteristics for separating of bifurcating *Astragalus* (Fabaceae) on sectional level. *Turkish Journal of Botany* 28(1/2): 241–245.
- Ghahremaninejad, F., Joharchi, M.R., Fereidounfar, S. & Hoseini, E. 2016. *Astragalus orientopersicus*, sp. nov. (Fabaceae), a new taxon from the Khorassan province (Iran). *Adansonia* 38(1): 29–33.
- Ghahremaninejad, F., Joharchi, M.R. & Memariani, F. 2022. *Astragalus durandianus* (sect. *Trachycercis*) as an Iranian endemic species: emended description, distribution, and conservation. *Phytotaxa* 550(1): 32–44.
- Kazempour Osaloo, S., Maassoumi, A.A. & Murakami, N. 2003. Molecular systematics of the genus *Astragalus* L. (Fabaceae): Phylogenetic analyses of nuclear ribosomal DNA internal transcribed spacers and chloroplast gene *ndh F* sequences. *Plant Systematics and Evolution* 242(4): 1–32.
- Kazempour Osaloo, S., Maassoumi, A.A. & Murakami, N. 2005. Molecular systematics of the Old World *Astragalus* (Fabaceae) as inferred from nrDNA ITS sequence data. *Brittonia* 57(4): 367–381.
- Khan, A., Ahmad, M., Zafar, S., Abbas, Q., Arfan, M., Zafar, Sultana, S., Ullah, S.A., Khan, S., Akhtar, A., Kilic, O. & Ozdemir, F.A. 2022. Light and scanning electron microscopic observation of palynological characteristics in spineless *Astragalus* L. (Fabaceae) and its taxonomic significance. *Microscopy Research and Technique* 85(7): 2409–2427.
- Khan, S., Jan, G., Ahmad, M., Gul, F., Zafar, M., Mangi, J. Din, Bibi, H., Sultana, S., Usma, A. & Majeed, S. 2021. Morpho-palynological assessment of some species of family Asteraceae and Lamiaceae of District Bannu, Pakistan on the bases of light microscope & scanning electron microscopy. *Microscopy Research and Technique* 84(6): 1220–1232.
- Lashin, G.M. 2006. Comparative morphology of pollen grains of some taxa of tribe Trifolieae (Fabaceae: Papilionoideae) from Egypt. *International Journal of Botany* 2(3): 270–277.
- Maassoumi, A.A. 2005. The Genus *Astragalus* in Iran, Vol. 5. Research Institute of Forests and Rangelands, Tehran (In Persian).
- Maassoumi, A.A. 2020. The comprehensive online resource for Mega genus of *Astragalus*. Ver. 1 (Published in the Internet).
- Mahmoodi, M., Ghahremaninejad, F. & Maassoumi, A.A. 2022. Diversity of vascular plants in Damirli Mountains (Zanjan Province, NW of Iran). *Rostaniha* 23(Suppl. 1): 1–131 (In Persian).
- Majeed, S., Zafar, M., Ahmad, M., Kilic, O., Sultana, S., Raza, J., Yaseen, G., Gul, H., Mir, S., Lubna & Jabeen, M. 2020. Pollen morphological investigations of family Cactaceae and its taxonomic implication light microscopy and scanning electron microscopy. *Microscopy Research and Technique* 83(7): 767–777.
- Mirzaei, L., Assadi, M., Nejadstari, T. & Mehregan, I. 2015. Comparative seed and leaf micromorphology of *Colutea* species (Fabaceae) from Iran. *Environmental and Experimental Biology* 13(4): 183–187.
- Nabila, Ahmad, M., Zafar, M., Bahadur, S., Sultana, S., Taj, S., Celep, F., Majeed, S. & Rozina, 2022.

- Palynomorphological diversity among the Asteraceous honeybee flora: An aid to the correct taxonomic identification using multiple microscopic techniques. *Microscopy Research and Technique* 85(2): 570–590.
- Noroozi, J., Talebi, A., Doostmohammadi, M. & Bagheri, A. 2020. The Zagros Mountain Range. *Plant Biogeography and Vegetation of High Mountains of Central and South-West Asia*. Pp. 185–214.
- Oskouian, R., Kazempour Osaloo, S. & Maassoumi, A.A. 2007. Pollen morphology of *Astragalus* section *Malacothrix* and its allies (Fabaceae) in Iran. *Journal of Science (University of Tehran)* 32(4): 177–181.
- Osman, A., Al-Ghamdi, F. & Guetat, A. 2014. Pollen morphology of some species of genus *Astragalus* L. (Fabaceae) in Northern region of Saudi Arabia. *Life Science Journal* 11(11): 1006–1019.
- Ozbek, F., Ekici, M., Buyukkartal, H. & Pinar, N. 2021. Anatomy, palynology and micromorphology of the genus *Astragalus* L. (Fabaceae) section *Uliginosi* Gray in Turkey. *Journal of the Institute of Science and Technology* 11(4): 2525–2536.
- Perveen, A. & Qaiser, M. 1998. Pollen flora of Pakistan-VIII Leguminosae (subfamily: Papilionoideae). *Turkish Journal of Botany* 22(2): 73–92.
- Pinar, M., Ekici, M., Aytac, Z., Akan, H., Ceter, T. & Alan, S. 2009. Pollen morphology of *Astragalus* L. sect. *Onobrychoidei* DC. (Fabaceae) in Turkey. *Turkish Journal of Botany* 33(4): 291–303.
- Podlech, D. & Zarre, Sh. (with collaboration of Ekici, M., Maassoumi, A.A. & Sytin, A.) 2013. A Taxonomic Revision of the Genus *Astragalus* L. (Leguminosae) in the Old World. Vols 1–3. *Naturhistorisches Museum, Wien*.
- Ranjbar, M., Assadi, A. & Karamian, R. 2012. Notes on *Astragalus* sect. *Macrophyllum* with a Cytogenetic Report on its two Tetraploid Species. *Journal of Science (Tarbiat Moallem University)* 11(2): 211–226.
- Raza, J., Ahmad, M., Zafar, M., Athar, M., Sultana, S., Majeed, S., Yaseen, G., Imran, M., Nazish, M. & Hussain, A. 2020. Comparative foliar anatomical and pollen morphological studies of Acanthaceae using light microscope and scanning electron microscope for effective microteaching in community. *Microscopy Research and Technique* 83(9): 1103–1117.
- Ullah, F., Zafar, M., Ahmad, M., Dilbar, S., Shah, N., Sohail, A., Zaman, W., Iqbal, M., Bahadur, S. & Tariq, A. 2018. Pollen morphology of subfamily Caryophylloideae (Caryophyllaceae) and its taxonomic significance. *Microscopy Research and Technique* 81(7): 704–715.
- Umer, F., Zafar, M., Ullah, R., Bari, A., Khan, M.Y., Ahmad, M. & Sultana, S. 2022. Copy for pollen morphology of selected taxa of family Asteraceae and Brassicaceae. *Microscopy Research and Technique* 85(1): 373–384.
- Uzun, A., Aytac, Z. & Tulucu, F. 2021. *Astragalus nurhakedagensis* (sect. *Hololeuce* Bunge /Fabaceae), a new species from Turkey. *Turkish Journal of Botany* 45(6): 573–586.
- Zarre, S. 2003. Hair micromorphology and its phylogenetic application in thorny species of *Astragalus* (Fabaceae). *Botanical Journal of the Linnean Society* 143(3): 323–330.
- Zarre, S. & Azani, N. 2013. Perspectives in taxonomy and phylogeny of the genus *Astragalus* (Fabaceae): a review. *Progress in Biological Sciences* 3(1): 1–6.