

Original Article

Prevalence and Risk Factors of Various Gastrointestinal Malignancies

Akram Ahangarpour^{1,2}, Roya Salehi Kahyesh^{3*}, Fereashteh Nemati³, Mohamad Hosein Rastegar³, Saeid Bitaraf³, Vahid Radmehr²

1. Department of Physiology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

2. Diabetes Research Center, Health Research Institute, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

3. Thalassemia & Hemoglobinopathy Research Center, Health Research Institute, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

Article Info:

Received: 11 August 2024

Revised: 23 September 2024

Accepted: 28 September 2024

Keywords:

Prevalence, Risk Factors,
Gastrointestinal Tract
Malignancy.

ABSTRACT

Gastrointestinal (GI) cancers are among the most common and fatal tumors worldwide, with incidence rates varying significantly due to factors such as inheritance, lifestyle, and diet. Understanding the prevalence and related risk factors is essential for improved prevention and treatment techniques. Data were collected from patients admitted to Baqaei Hospital during 2019 to 2022. The questionnaire included demographic data (age and sex), occupation, location of malignancy, genetic history, smoking habits, underlying disease (hypertension, diabetes, rheumatism), performing physical activities (exercise), presence of metastasis, and complete survival status. The data analysis of this study was performed using statistical software: SPSS. Both linear regression and unadjusted logistic regression analyses were conducted. T-test and chi-square statistical methods were also used for analysis in this study. The results revealed that the survival rate in patients with gastrointestinal malignancies are significantly associated with factors such as age, family history, tumor location (colon and pancreas), history of smoking, concomitant disease, metastasis (rectal cancer), and physical activity. The average age of death is 66.05 years. Among them, approximately 20.4% had a family history of cancer. Also, 50% exhibited metastasis, 44.8% had a history of smoking, 74.8% had underlying diseases, and 99.6% had not exercised. Physical activity was significantly lower in deceased patients, and fewer deaths occurred among those with high physical activity levels. Rectal cancer had the highest percentage of metastasis among living and deceased patients. It will be useful to carry out more studies to determine the clinical and demographic factors that affect the survival of patients with colorectal cancer patients. Public awareness and early medical consultation are crucial for improving outcomes.

Corresponding Author:

royaarta@yahoo.com



<https://orcid.org/0000-0002-0601-6842>

<https://orcid.org/0000-0003-4688-9856>

How to cite this article: Ahangarpour A, Salehi Kahyesh R, Nemati F, Rastegar MH, Bitaraf S, Radmehr V. Prevalence and Risk Factors of Various Gastrointestinal Malignancies. *Archives of Razi Institute*. 2025;80(3):719-724. DOI: 10.32592/ARI.2025.80.3.719



1. Introduction

Gastrointestinal cancer involves a large number of cases that often leads to death. Metastases from the esophagus, stomach, liver, pancreas, gall bladder, colon, and rectum are the most prevalent locations of gastrointestinal cancer (1). These cancers account for approximately 37% of all cancer-related deaths worldwide (2). In Iran, gastrointestinal cancers are highly prevalent, with 20,719 deaths, comprising 44.4% of all cancer-related mortalities. Stomach cancer is the most common cancer among men in Iran and ranks third after breast and colorectal cancer in women (3). It is the leading cause of death from gastrointestinal cancers, followed by esophageal and colorectal neoplasms (4).

Factors contributing to the development of gastrointestinal cancers include dietary habits such as salty foods and processed foods, low consumption of vegetables and fruits, a sedentary lifestyle, smoking, alcohol use, advanced age, male gender, family history of cancer, and *Helicobacter pylori* infection (5, 6).

Also, rapid industrialization and urbanization, particularly in large cities, have increased pollution levels, further elevating cancer risks (5). Finally, all these factors increase the incidence of gastrointestinal cancers (7). These cancers are particularly significant due to their prevalence, high mortality rates, and their prominence among other cancer types (5). The challenges associated with the treatment of many types of cancer, along with the high costs involved, have placed a significant burden on national health budgets (5). Early and timely diagnosis of gastrointestinal cancers, particularly through endoscopy, is crucial as it can significantly improve patient survival rates and reduce treatment costs (8-10). Given the increasing incidence and mortality rates associated with gastrointestinal cancers, this study aims to investigate the prevalence and risk factors of various gastrointestinal malignancies in patients at Baqaei 2 Hospital in Ahvaz.

2. Materials and Methods

2.1. Data Collection

In this retrospective study, data related to primary gastrointestinal tract cancers were obtained from the archived patient files at the Health Technology Unit of Baqaei Hospital 2, covering four years from 2019 to 2022.

2.2. Methodology

The questionnaire collected comprehensive patient information, including demographic data (age and sex), occupation, location of malignancy, family history of

cancer, tobacco use (smoking and other types), alcohol consumption, underlying conditions (such as hypertension, diabetes, and rheumatism), physical activity (exercise), presence of metastasis, and survival status (alive/deceased). The collected data were then analyzed statistically.

2.3. Statistical Analysis

2.3.1. Statistical software

SPSS, linear regression, and unadjusted logistic regression were used for data calculation and analysis. Also, the study's participants were described using the mean and standard deviation in the presence of continuous data, the number and percentage in the presence of classified data. T-test and chi-square statistical methods were also used for analysis in this study. Significance levels were considered 0.05.

3. Results

In this four-year study, data from 999 patients with various types of gastrointestinal malignancies were analyzed. The results indicated that the average age of deceased patients was 66.05 years, while the average age of surviving patients was 56.29 years, with a significant difference between the two groups ($P < 0.05$). Among the patients, 432 were female (43.3%) and 567 were male (56.7%).

Further analysis revealed a significant relationship between patient survival and factors such as family history, smoking, underlying disease, metastasis, physical activity, occupation and tumor location ($P < 0.05$). Additionally, survival rates were lower in patients with a positive family history, smoking habits, underlying disease, metastasis, lack of physical activity, and malignancies identified in the colon and pancreas (Table 1).

An analysis of the predictors of mortality in individuals with gastrointestinal cancer revealed that age, tumor location, family history of cancer, smoking habits, underlying diseases, and metastasis are significantly associated with cancer-related death. This study also found that the presence of gastrointestinal cancers in first- and second-degree relatives plays a crucial role in the development of these malignancies (Table 2).

Metastases to the esophagus, stomach, colon, bones, liver, and lungs were associated with higher mortality in patients, whereas metastases to the kidneys were associated with the lowest death rate.

Table 1. Baseline characteristics of participants.

Variable		alive	dead	p-value
		N=729	N=270	
age		56.29 (14.42)	66.05 (15.62)	<0.001
sex	female	327 (44.9%)	105 (38.9%)	0.098
	male	402 (55.1%)	165 (61.1%)	
Family history	negative	676 (92.7%)	215 (79.6%)	<0.001
	positive	53 (7.3%)	55 (20.4%)	
smoking	no	612 (84.0%)	149 (55.2%)	<0.001
	yes	117 (16.0%)	121 (44.8%)	
background disorder	no	528 (72.4%)	68 (25.2%)	<0.001
	yes	201 (27.6%)	202 (74.8%)	
metastasis	negative	686 (94.1%)	135 (50.0%)	<0.001
	positive	43 (5.9%)	135 (50.0%)	
exercise	no	698 (95.7%)	269 (99.6%)	<0.001
	yes	31 (4.3%)	1 (0.4%)	
occupation	disable	1(0.14)	10(3.70)	<0.001
	freelance	118(16.19)	37(13.70)	
	retired	109(14.95)	45(16.67)	
	jobless	151(20.71)	72(26.67)	
	housekeeper	310(42.52)	96(35.56)	
	studying	7(0.96)	1(0.37)	
tumor place	employee	33(4.53)	9(3.33)	<0.001
	appendix	3(0.41%)	1(0.37%)	
	pancreas	29(3.98%)	78(28.89%)	
	small intestine	1(0.14%)	0(0.00%)	
	rectum	56(7.68%)	12(4.44%)	
	esophagus	60(8.23%)	16(5.93%)	
	gastric	209(28.67%)	45(16.67%)	
	neuroendocrine	0(0.00%)	1(0.37%)	
	hepatoblastoma	8(1.10%)	2(0.74%)	
	liver	43(5.90%)	20(7.41%)	
	colon	299(41.02%)	90(33.33%)	
	gallbladder	21(2.88)	5(1.85)	

Table 2. Family history of cancer in first and second-generation.

Family history		Live situation	
	Dead (%)	Alive (%)	Total (%)
Negative	215(79.63)	676(92.73)	891(89.19)
pancreas first	1(0.37)	0(0)	1(0.10)
breast first	10(3.70)	12(1.65)	22(2.20)
breast second	0(0)	1(0.14)	1(0.10)
prostate first	2(0.74)	4(0.55)	6(0.60)
ovary first	10(3.70)	4(0.55)	14(1.40)
uterus first	1(0.37)	1(0.14)	2(0.20)
rectum first	2(0.74)	0(0)	2(0.20)
lung first	1(0.37)	1(0.14)	2(0.20)
leukemia first	2(0.74)	2(0.27)	4(0.40)
esophagus first	6(2.22)	5(0.69)	11(1.10)
esophagus second	0(0)	1(0.14)	1(0.10)
gastric first	9(3.33)	12(1.65)	21(2.10)
liver first	2(0.74)	0(0)	2(0.20)
colon first	9(3.33)	9(1.23)	18(1.80)
gallbladder first	0(0)	1(0.14)	1(0.10)
Total	270	729	999
	100.00	100.00	100.00

4. Discussion

Considering the high prevalence of gastrointestinal cancers in Iran, particularly esophageal and stomach cancer, this study aimed to investigate the prevalence and risk factors of various types of gastrointestinal malignancies in patients at Baqaei 2 Hospital. The findings showed that the survival rate across all types of gastrointestinal cancers significantly correlated with age, with higher mortality observed in older patients. A study by Ebrahimi et al. in 2024, which investigated the mortality trends of gastrointestinal cancers in Babol, Northern Iran (2011-2013), also showed that the mortality rate in gastrointestinal cancers increases with age (11). Similarly, Salehzadeh et al., (2018) in a study titled "The Annual Mortality Trend of Gastrointestinal Cancers in Iran during the Years 1990 to 2015," used data from the Iranian Civil Registration System (1995 to 2010) and two cemeteries in Tehran (1995 to 2010) and Isfahan (2007 to 2010). The study found that cancer mortality from all causes increased with age and was more prominent in adults aged 50 years or beyond (8), consistent with the results of our study, where the mortality rate of most malignancies increases with age.

In the present study, a comparison of survival rates among patients with gastrointestinal malignancies based on family history revealed a significant relationship between survival rate and family history. Similarly, Ghasemi et al. (2023) reported a significant relationship between patient survival and a family history of cancer (12).

Yousefi et al. (2018), in a review study titled "Risk Factors for Gastric Cancer," in 2017, performed a structured overview using databases such as Science Direct, Scopus, PubMed, Cochrane, and Web of Science (ISI). This study identified and categorized 52 risk factors for stomach cancer into nine major categories: diet, lifestyle, genetic predisposition, family history, medical treatments and conditions, infections, demographic characteristics, occupational exposures, and ionizing radiation (13). These findings support our observations.

Regarding smoking, our analysis revealed a significant inverse relationship between smoking history and patient survival. Chen et al. (2024) conducted a study on the prevalence, types, and risk factors of gastrointestinal tract diseases in Hainan Province, China. They concluded that smoking significantly reduces the survival rate of patients with gastrointestinal malignancies (14). Similarly, Wong

et al. (2019) reported that non-modifiable factors such as genetic predisposition, ethnicity, age, gender, family history, smoking habits, alcohol consumption, weight (BMI), Western diet, low physical activity, chronic diseases, and microbiota influence the prevalence and risk factors of colorectal cancer in Asia (15), which is consistent with our findings. Although the exact mechanism linking smoking to gastrointestinal symptoms is not fully understood, previous studies have established a connection between smoking and various gastrointestinal disorders, including gastroesophageal reflux disease, esophageal cancer, gastric ulcers, and gastric cancer (16).

Our study also indicated a significant relationship between survival rates and the presence of underlying diseases and comorbidities, with a notably higher percentage of comorbidities observed in deceased patients. A 2021 study titled "Prevalence and Risk Factors of Upper Gastrointestinal Cancers during Endoscopy" cancers diagnosed less than 6 months after endoscopy were considered "common". While those diagnosed between 6 and 36 months were categorized as "missed". The study found that esophageal adenocarcinomas were missed more frequently than squamous cell cancers (6.1% vs. 4.2%), with a relative risk of 1.4. Additionally, most gastric cancers were adenocarcinomas, with 5.7% classified as missing. This study identified comorbidities as a significant risk factor for gastrointestinal malignancies (17).

In this research, the survival rate of patients with gastrointestinal malignancies was significantly related to the presence of metastasis, with a notably higher percentage of positive metastases observed in deceased patients. Various studies have highlighted that tumor size and the number of involved lymph nodes are critical factors, affecting patient survival. As tumor size and the number of involved lymph nodes increase, the likelihood of metastasis rises, ultimately reducing patient survival rates (18, 19). These findings highlight the importance of early detection and treatment, as patients who are unaware of their disease in its initial stages are more likely to experience disease progression and metastasis. Effective strategies to prevent metastasis are crucial for improving patient outcomes.

The findings of the present study indicated a significant relationship between survival rates of patients with gastrointestinal malignancies and physical activity. Deceased patients exhibited notably lower levels of

physical activity. Vishwanath et al. (2024) reported that lifestyle, genetics, and environmental factors contribute significantly to the rising incidence of gastrointestinal malignancies among younger populations, since physical activity is associated with increased survival rates in these patients (20).

In our study, examining the survival rates of patients with various types of gastrointestinal malignancies based on the primary tumor location revealed a significant relationship between these variables. In both patient groups, colon tumors were the most prevalent, while pancreatic cancer was more significantly associated with deceased patients. Ebrahimi et al. (2024) observed a significant relationship between tumor location and survival, with the highest rates of malignancy found in the colon and pancreas, which notably affected survival rates (11).

Additionally, Alhazmi et al. (2020) conducted a case-control study titled "Prevalence and Patterns of Gastrointestinal Cancers in Obese Patients" at a teaching hospital in Saudi Arabia. Using medical records of adult patients diagnosed with gastrointestinal cancer from January 2010 to May 2018 at King Abdulaziz University Hospital, the study included 532 patients. It found that colorectal cancer was the most common tumor site in obese patients, followed by gastric and pancreatic cancers (21), which is consistent with the findings of the present study. Finally, the distribution of metastasis among the patients in the present study showed that the highest percentage of metastasis, both in deceased and surviving patients, were in the rectum. Based on the study by Rosenberg et al. rectal tumors have a worse prognosis than colon tumors (22), which is consistent with the present study.

The results of this present study indicated that survival rates in patients with gastrointestinal malignancies are significantly related to several factors, including age, family history, tumor location (particularly colon and pancreas), smoking history, comorbidities, metastasis (notably in rectal cancer), and physical activity. Rectal cancer was exhibited the highest percentage of metastasis among both living and deceased patients. To better understand the clinical and demographic factors affecting the survival of patients with colorectal cancer, further research is warranted. Additionally, public awareness to promote early medical consultation and timely diagnostic procedures is essential.

Acknowledgment

This article is a result of Ms. Fereshteh Nemati's general medicine thesis. This research was supported by financial support (U-02460) from Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. We thank the dedicated laboratory team of Baghaei 2 Hospital for their sincere cooperation.

Authors' Contribution

Study concept and design, administration of technical and material support, and study supervision: R. SK, A. A.

Data acquisition: M. HR, F. N.

Analysis and interpretation of data, statistical analysis: S. B.

Drafting of the manuscript and critical revision for important intellectual content: V. R, R. SK.

Ethics

This study was approved by the ethics committee under the code IR.AJUMS.REC.1402.086, at Jundishapur University of Ahvaz.

Conflict of Interest

The authors declare no conflicts of interest related to this study.

Data Availability

The data supporting the findings of this study are available on request from the corresponding author.

References

1. Xie Y, Shi L, He X, Luo Y. Gastrointestinal cancers in China, the USA, and Europe. *Gastroenterology report*. 2021;9(2):91-104.
2. Huang J, Lucero-Prisno III DE, Zhang L, Xu W, Wong SH, Ng SC, et al. Updated epidemiology of gastrointestinal cancers in East Asia. *Nature Reviews Gastroenterology & Hepatology*. 2023;20(5):271-87.
3. Swain CK. Environmental pollution indices: a review on concentration of heavy metals in air, water, and soil near industrialization and urbanisation. *Discover Environment*. 2024;2(1):5.
4. Zarea K, Beiranvand S, Ghanbari S, Tuveson H. Incidence of gastrointestinal cancers in Iran: a systematic review. *Jundishapur J Chronic Dis Care*. 2017;6(1):e37224.
5. Lu L, Mullins CS, Schafmayer C, Zeißig S, Linnebacher M. A global assessment of recent trends in

gastrointestinal cancer and lifestyle-associated risk factors. *Cancer Communications*. 2021;41(11):1137-51.

6. Jardim SR, de Souza LMP, de Souza HSP. The rise of gastrointestinal cancers as a global phenomenon: unhealthy behavior or progress? *International journal of environmental research and public health*. 2023;20(4):3640.

7. Li P, Jing J, Guo W, Guo X, Hu W, Qi X, et al. The associations of air pollution and socioeconomic factors with esophageal cancer in China based on a spatiotemporal analysis. *Environmental Research*. 2021;196:110415.

8. Salimzadeh H, Delavari F, Sauvaget C, Rezaee N, Delavari A, Kompani F, et al. Annual trends of gastrointestinal cancers mortality in Iran during 1990-2015; NASBOD study. *Archives of Iranian medicine*. 2018;21(2):46-55.

9. Nijjar GS, Aulakh SK, Singh R, Chandi SK, kaur Aulakh S. Emerging Technologies in Endoscopy for Gastrointestinal Neoplasms: A Comprehensive Overview. *Cureus*. 2024;16(6).

10. Ding Y, Ren L, Geng Y, Fu C, Shi R. The current status and prospects of early diagnosis and treatment of esophageal cancer in China. *Cancer Screening and Prevention*. 2024;3(2):106-12.

11. Ebrahimi P, Karami M, Delavari S, Shojaie L, Hosseini-Berneti S-H, Bayani F, et al. Investigating the mortality trend of gastrointestinal cancers in Babol, North Iran (2013–2021). *BMC gastroenterology*. 2024;24(1):100.

12. Ghasemi SS, Zarchi AAK, Alimohamadi Y, Raei M, Sepandi M. Survival analysis and its related factors among patients with breast cancer referred to a military hospital in Tehran. *EBNESINA*. 2023;25(1):4-12.

13. Yusefi AR, Lankarani KB, Bastani P, Radinmanesh M, Kavosi Z. Risk factors for gastric cancer: a systematic review. *Asian Pacific journal of cancer prevention: APJCP*. 2018;19(3):591.

14. Chen C, Zhang D-y, Chen S, Huang S, Zeng F, Li D, et al. Prevalence, types, and risk factors of functional gastrointestinal diseases in Hainan Province, China. *Scientific reports*. 2024;14(1):4553.

15. Wong MC, Ding H, Wang J, Chan PS, Huang J. Prevalence and risk factors of colorectal cancer in Asia. *Intestinal research*. 2019;17(3):317-29.

16. Yuan S, Chen J, Ruan X, Sun Y, Zhang K, Wang X, et al. Smoking, alcohol consumption, and 24 gastrointestinal diseases: Mendelian randomization analysis. *elife*. 2023;12:e84051.

17. Januszewicz W, Witczak K, Wieszczy P, Socha M, Turkot MH, Wojciechowska U, et al. Prevalence and risk factors of upper gastrointestinal cancers missed during endoscopy: a nationwide registry-based study. *Endoscopy*. 2022;54(07):653-60.

18. Karimi A, Delpisheh A, Sayehmiri K, Saboori H, Rahimi E. Predictive factors of survival time of breast cancer in kurdistan province of Iran between 2006-2014: a cox regression approach. *Asian Pacific Journal of Cancer Prevention*. 2014;15(19):8483-8.

19. Hajihosseini M, Faradmal J, Sadighi-Pashaki A. Survival analysis of breast cancer patients after surgery with an intermediate event: application of illness-death model. *Iranian Journal of Public Health*. 2015;44(12):1677.

20. Vishwanath A, Krishna S, Manudhane AP, Hart PA, Krishna SG. Early-onset gastrointestinal malignancies: An investigation into a rising concern. *Cancers*. 2024;16(8):1553.

21. Ahangarpour A, Salehi Kahyesh R, Nemati F, Rastegar MH, Bitaraf S, Radmehr V. Prevalence and risk factors of various gastrointestinal malignancies. *Archives of Razi Institute*. 2024.

22. Rebersek M. Gut microbiome and its role in colorectal cancer. *BMC cancer*. 2021;21(1):1325.