

Short Communication

Investigation of Antibiotic Susceptibility Patterns in Bacteria Isolated During a Male Sheep Castration Surgery

Zahra Akbari¹, Neli Zalikani¹, Mohaddeseh Babaei^{1*}

1. Department of Veterinary Medicine, Babol Branch, Islamic Azad University, Babol, Iran

Article Info:

Received: 22 September 2024

Revised: 3 January 2025

Accepted: 16 January 2025

Keywords:

Ampicillin,
Antibiotic susceptibility,
Bacterial identification,
Drug resistance, *Yersinia* spp.

ABSTRACT

There are different antibiotic resistance profiles among *Yersinia* spp. This pathogenic bacterial species causes yersiniosis worldwide, requiring testing of the organism's susceptibility in the local environment. An antibiotic susceptibility profile of a *Yersinia* spp. isolated from a castration surgical site was analyzed to provide insight into selecting appropriate antibiotics to treat *Yersinia* spp. infections while addressing antibiotic resistance issues effectively. The surgical site was swabbed before castration, and cultures were performed. Samples from the surgical site were taken after the procedure, cultured, and then incubated. CLSI 2020 guidelines were followed for interpreting antibiotic susceptibility tests. The Kirby-Bauer disk diffusion method was applied to better understand antibiotic susceptibility and resistance patterns, and a zone of inhibition measurement was used to determine the antibiotic effectiveness. Staining and microscopic examination of swab samples after surgery revealed a single colony of Gram-negative bacteria. Laboratory tests confirmed that the isolated Gram-negative bacilli were indeed *Yersinia* spp. Methyl Red and Voges-Proskauer tests showed negative results, while Citrate utilization testing demonstrated a positive result. A positive result was obtained for *Yersinia* spp. in the glucose fermentation test. Specifically, nitrofurantoin showed a significant zone of inhibition of over 17 mm, and gentamicin showed a zone of inhibition greater than 27 mm. However, resistance to ampicillin (11 mm), ceftriaxone, and cefazolin was observed. Due to the observed resistance to antibiotics, our results indicate that nitrofurantoin and gentamicin are likely to be the best options for treating *Yersinia* spp., in contrast to ampicillin, cefazolin, and ceftriaxone, which may be unsuitable due to resistance.

Corresponding Author:

mohadesebabaei9973@gmail.com

<https://orcid.org/0009-0004-3687-2852>

<https://orcid.org/0009-0003-4778-6151>

How to cite this article: Akbari Z, Zalikani N, Babaei M. Investigation of Antibiotic Susceptibility Patterns in Bacteria Isolated During a Male Sheep Castration Surgery. *Archives of Razi Institute*. 2025;80(4):1075-1080. DOI: 10.32592/ARI.2025.80.4.1075



1. Introduction

There is no doubt that antibiotic resistance is one of the most pressing public health crises facing both the human and animal populations in the modern world. *Yersinia* spp., specifically *Yersinia enterocolitica*, is an example of an organism that develops resistance to a wide range of antibiotics (1, 2). The extensive and indiscriminate application of antibiotics in veterinary medicine contributes to antibiotic resistance. Despite the inherent risks of bacterial contamination, it is still common practice to use these substances in surgical procedures aimed at sterilizing animals (3, 4). The emergence of *Yersinia*'s antibiotic-resistant strains has substantial consequences beyond geographic boundaries (5).

These resilient strains challenge the effectiveness of treatment methods and animal health practices. Addressing antibiotic resistance in *Yersinia* spp., and related pathogens is imperative due to animal and human health interdependence. The problem of bacterial resistance in veterinary medicine can further complicate public health concerns. It is vital to investigate *Yersinia* spp. susceptibility patterns to a wide range of antibiotics. A critical step towards safeguarding public health is understanding how bacteria respond to various antibiotics. The current study aims to determine which antibiotics are effective against *Yersinia* spp. when they adapt to antibiotics. This study includes insights into how veterinary medicine can make better decisions regarding treatment during sterilization procedures for better outcomes.

2. Materials and Methods

2.1. Sampling

Samples were collected from the surgical site before preparation for castration surgery in a sheep. Before starting the surgery, the site was thoroughly cleaned with alcohol, chlorhexidine, and povidone-iodine. The surgical site and suture location were swabbed following ceftriaxone administration and the completion of the procedure. After 24 hours of incubation of samples, following linear culture, crystal violet, iodine, and fuchsin stains were applied to determine the colony type.

As soon as bacteria were identified through Gram staining, they were cultured linearly within their specific media. To cultivate Gram-negative bacteria, including *Yersinia*, the EMB medium was used, along with Simmon's citrate, TCI, SIM, and MR-VP. A 24-hour

incubation at 37°C was used to promote bacterial growth and stability after being cultured in their specialized media. Testing with Methyl Red was performed using an MR medium. A VP test was conducted using an MRVP medium divided into two parts. SIM medium, which is semisolid, is then used for further investigation (6).

2.2. Antibiotic susceptibility test

Gram-negative bacteria were cultured on Mueller-Hinton agar medium once removed from the refrigerator. By inserting six antibiotic disks, this study examined the susceptibility of several antibiotics, including nitrofurantoin, gentamicin, cefazolin, tetracycline, ceftriaxone, and ampicillin. The antibiotics chosen for this study were relevant to veterinary medicine, since veterinarians may use them postoperatively. Sterilization was achieved by disinfecting the disks with alcohol and incubating them in the culture medium. At 37°C, the plates were incubated for 24 hours. The inhibition zones were measured with a ruler after 24 hours of incubation. Antibiotic susceptibility testing was performed using standards established by the Clinical and Laboratory Standards Institute (CLSI) (7, 8).

3. Results

3.1. Sampling culture findings

Among the colonies sampled before surgery, Gram-positive cocci, Gram-positive bacilli, and Gram-negative bacilli were found. After surgery, one Gram-negative colony was found in swab samples after staining and microscopic examination. An extensive series of biochemical tests determined the bacteria's identity. Testing the bacteria with the Methyl Red (MR) test determined that they did not produce acidic end products when they metabolized glucose. There was no evidence of acetoin production by the bacteria in Voges-Proskauer tests (VP). This suggests that enteric bacteria do not produce acetoin. During the Citrate utilization test, it was determined that the bacterium could utilize citrate alone as a carbon source. The bacteria showed positive results in a glucose fermentation test, indicating they can ferment glucose. A lack of these components prevented them from producing hydrogen sulfide (H₂S), indole, or exhibiting motility. These tests contributed to identifying *Yersinia* spp. as the pathogen in the case under investigation.

3.2. Antibigram

The inhibition zones for each antibiotic were measured to determine the bacterium's susceptibility to antibiotics (Figure 1).



Figure 1. The antibiotic disks were inserted into the media, allowing six different antibiotics to be tested for *Yersinia* resistance. The antibiotics utilized were ceftriaxone, nitrofurantoin, gentamicin, cefazolin, ampicillin and Tetracycline.

In addition, this bacterium displayed an extensive inhibitory zone (more than 17 mm), which qualified it as highly sensitive to nitrofurantoin. There was also evidence to indicate that the bacteria are highly susceptible to gentamicin, based on the fact that the zone diameter was 27 mm. Furthermore, tetracycline also showed an effect against the bacteria, displaying a zone diameter of 23 mm, indicating susceptibility. The zone diameter of the bacterium was only 11 mm, despite its resistance to ampicillin. As the zone diameters were 17 mm and 10 mm, it was evident that the bacterium was resistant to ceftriaxone and cefazolin (Table 1).

4. Discussion

One of the critical pathogenic bacteria is *Yersinia* spp., a member of the Enterobacteriaceae family (9). There are several serotypes and biotypes of this bacterium. Among the factors contributing to its pathogenicity are its ability to grow at temperatures between 0 and 44°C and the diversity of its surface antigens. *Yersinia* spp. causes yersiniosis, a zoonotic disease affecting both humans and animals (10). Antibiotics are used in veterinary medicine as a treatments, preventatives, and even growth promoters, to manage infections and prevent infectious diseases (11).

Table 1. Analyses of antibiotic sensitivity and resistance in *Yersinia* spp.

Antibiotic	μAmount (g)	Zone Size (mm)	Sensitive	Intermediate	Resistance
Nitrofurantoin	300	28	17≤	16-15	14≥
Gentamicin	10	27	15≤	13-14	12≥
Tetracycline	30	23	15≤	14-12	11≥
Ceftriaxone	30	17	23≤	22-20	19≥
Ampicillin	10	11	17≤	16-14	13≥
Cefazolin	30	10	23≤	22-20	19≥

However, resistance to antibiotics is on the rise, affecting both bacterial populations and different hosts. Evidence suggests that horizontal gene transfer and mobile genetic elements are responsible in the development of this resistance, which can reduce the effectiveness of antimicrobial agents in both humans and animals (12).

According to a study conducted in Bulgaria, *Yersinia* spp., isolated from pork were resistant to ampicillin, tetracycline, and nalidixic acid, yet sensitive to chloramphenicol and gentamicin (13). In Egypt, another study found the highest resistance to ampicillin, cefazolin, and amoxicillin/clavulanic acid among strains of *Yersinia* spp (14). Numerous studies have documented diverse antibiotic resistance patterns in *Yersinia* isolates (15, 16). As a result of these patterns, it is imperative conducting localized susceptibility testing in specific regions (17). To successfully treat bacterial infections, it is crucial to choose effective antibiotics, particularly for highly pathogenic bacteria such as *Yersinia*.

Based on the results of the current study, the isolate of *Yersinia* spp. from castration surgical sites was very susceptible to nitrofurantoin and gentamicin. Due to their close relationship, these antibiotics may effectively treat and eradicate *Yersinia* spp. infections. The treatment of *Yersinia* spp. infection should not involve antibiotics like ampicillin, ceftriaxone, or cefazolin due to bacterial

resistance to these drugs. Limitations of the present study include the small sample size and the limited number of control groups. In this study, we only examined bacteria isolated from castration surgery of male sheep, which may limit the generalizability of the results. To increase the accuracy and generalizability of the results, the number of samples in future studies should be increased, and the inclusion of additional control groups should be considered. Moreover, repeating the experiments under different conditions and using different laboratory methods can help improve the results' validity and accuracy. In addition, using animals from different breeds and conditions—especially in different geographical areas—can provide more information about antibiotic resistance patterns in various bacteria and help design more effective treatment strategies (Table 2).

As a revolutionary system, it is highly recommended that nanotechnology be considered for overcoming the challenge of drug resistance (18). Nanotechnology has attracted much attention due to its potential to improve disease diagnosis, treatment, and prevention, especially in human and veterinary medicine (19). Important applications of nanotechnology include designing nanoparticles for precise drug targeting, faster identification of pathogens, and the creation of intelligent drug delivery systems (20).

Table 2. This table can be a helpful tool to present suggestions and challenges that researchers can address in future studies, ensuring more comprehensive and valid results.

Challenge	Description	Suggested Improvement
Sample Size	The current study had a small sample size, limiting the ability to generalize findings.	Increase the sample size to ensure more robust and reliable results.
Control Group Limitations	Only a limited number of control groups were used, which may not reflect the broader spectrum of cases.	Add more control groups with diverse conditions to improve the validity of comparisons.
Replication of Experiments	The experiments were not repeated under different conditions, affecting the results' reproducibility.	Repeat experiments across different settings and methodologies to confirm consistency and reliability.
Animal Diversity	The study focused on a single breed of sheep, limiting the representation of different animal variations.	Include animals from different breeds and geographical regions to obtain a wider range of data on antibiotic resistance.
Geographical Variation	The study did not account for geographical differences in bacterial resistance patterns.	Perform studies in multiple regions to observe geographical trends in antibiotic resistance.
Data Interpretation	The limitations may influence the results of the testing methods and protocols used.	Use various testing methods to confirm findings and ensure comprehensive data interpretation.
Statistical Significance	The study might not have fully explored the statistical significance across different groups.	Perform deeper statistical analyses to explore the significance of observed differences in greater detail.

This technology can also help develop more accurate diagnostic tools and optimize treatments for drug-resistant diseases. Therefore, further research and investment in this area promise significant advances in various scientific and medical fields.

Various bacterial strains exhibit different antibiotic resistance profiles, so tailored treatment must be considered. According to the findings of this study, nitrofurantoin, Gentamicin, and Tetracycline may also be more effective than ampicillin, cefazolin, and ceftriaxone against *Yersinia* spp.

Acknowledgment

We sincerely thank Islamic Azad University, Babol Branch, Babol, Iran, for supporting us.

Authors' Contribution

Study concept and design: MB

Acquisition of data: ZA, NZ, MB

Analysis and interpretation of data: ZA, NZ, MB

Drafting of the manuscript: ZA, NZ, MB

Critical revision of the manuscript for important intellectual content:

Statistical analysis: ZA, NZ, MB

Administrative, technical, and material support: MB

Ethics

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed. IR.IAU.BABOL.REC.1403.065.

Conflict of Interest

The authors declare no conflict of interest.

Funding

No funding was received.

Data Availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author upon reasonable request.

References

- Shoaib M, Shehzad A, Raza H, Niazi S, Khan IM, Akhtar W, et al. A comprehensive review on the prevalence, pathogenesis and detection of *Yersinia enterocolitica*. RSC advances. 2019;9(70):41010-21
- Grudlewska-Buda K, Bauza-Kaszewska J, Wiktorczyk-Kapischke N, Budzyńska A, Gospodarek-Komkowska E, Skowron K. Antibiotic Resistance in Selected Emerging Bacterial Foodborne Pathogens—An Issue of Concern? Antibiotics. 2023;12(5):880
- Burgess BA. Prevention and surveillance of surgical infections: A review. Veterinary Surgery. 2019;48(3):284-90
- Saleh M, Harkinezhad MT, Marefat A, Salmani V. An outbreak of abortion in Afshari sheep with probable involvement of *Campylobacter* fetus. 2013
- Abdelwahab AM, El-Tawab A, Awad A, Abdallah F, Maarouf AA. Phenotypic and genotypic studies on antibiotic resistant *Yersinia enterocolitica* isolated from milk and milk products in Kaliobia, Egypt. Benha Veterinary Medical Journal. 2021;40(2):149-53
- Gita R, Kumaji SS, Lainjong EA. Identification of *Escherichia coli* bacteria in quail eggs that for sale in the central market of the city of Gorontalo. Journal of Health, Technology and Science (JHTS). 2021;2(2):19-27
- Humphries R, Bobenchik AM, Hindler JA, Schuetz AN. Overview of changes to the clinical and laboratory standards institute performance standards for antimicrobial susceptibility testing, M100. Journal of clinical microbiology. 2021;59(12):10.1128/jcm.00213-21
- Weinstein MP, Lewis JS. The clinical and laboratory standards institute subcommittee on antimicrobial susceptibility testing: background, organization, functions, and processes. Journal of clinical microbiology. 2020;58(3):10.1128/jcm.01864-19
- Zadernowska A, Chajęcka-Wierchowska W, Łaniewska-Trokenheim Ł. *Yersinia enterocolitica*: a dangerous, but often ignored, foodborne pathogen. Food Reviews International. 2014;30(1):53-70
- Dheyab ZS. Clinically Important *Yersinia*: Minireview. Journal of Medical Research and Health Sciences. 2022;5(10):2295-306
- Lü Z, Su X, Chen J, Qin M, Sheng H, Zhang Q, et al. Prevalence, bio-serotype, antibiotic susceptibility and genotype of *Yersinia enterocolitica* and other *Yersinia* species isolated from retail and processed meats in Shaanxi Province, China. LWT. 2022;168:113962
- Durofil A, Maddela NR, Naranjo RA, Radice M. Evidence on antimicrobial activity of essential oils and herbal extracts against *Yersinia enterocolitica*—A review. Food Bioscience. 2022;47:101712

13. Angelovska M, Zaharieva MM, Dimitrova LL, Dimova T, Gotova I, Urshev Z, et al. Prevalence, Genetic Homogeneity, and Antibiotic Resistance of Pathogenic *Yersinia enterocolitica* Strains Isolated from Slaughtered Pigs in Bulgaria. *Antibiotics*. 2023;12(4):716
14. Dowidar HA, Khalifa MI. Virulence Genotyping Profiles of Cefazolin Resistance *Yersinia enterocolitica* Isolated From Milk and Milk Products in Egypt. *Egyptian Journal of Veterinary Sciences*. 2023;54(4):555-69
15. Peng Z, Zou M, Li M, Liu D, Guan W, Hao Q, et al. Prevalence, antimicrobial resistance and phylogenetic characterization of *Yersinia enterocolitica* in retail poultry meat and swine feces in parts of China. *Food Control*. 2018;93:121-8
16. Younis G, Elkenany R, Dowidar H. Virulence genotyping and antimicrobial resistance profiles of *Yersinia enterocolitica* isolated from meat and meat products in Egypt. *Brazilian Journal of Biology*. 2020;81:424-36
17. Piras F, Spanu C, Sanna R, Siddi G, Mocci AM, Demontis M, et al. Detection, virulence genes and antimicrobial resistance of *Yersinia enterocolitica* in sheep and goat raw milk. *International Dairy Journal*. 2021;117:105011
18. Singh R, Smitha M, Singh SP. The role of nanotechnology in combating multi-drug resistant bacteria. *Journal of nanoscience and nanotechnology*. 2014;14(7):4745-56
19. Sadr S, Poorjafari Jafroodi P, Haratizadeh MJ, Ghasemi Z, Borji H, Hajjafari A. Current status of nano-vaccinology in veterinary medicine science. *Veterinary Medicine and Science*. 2023;9(5):2294-308
20. Talebi F, Misaghi A, Khanjari A, Kamkar A, Gandomi H, Saeedi M. Evaluation of antimicrobial activity of poly lactic acid (PLA) films containing cellulose nanoparticle and *Bunium persicum* and *Mentha piperita* essential oils (EOs). 2017