



Original Article

The First Molecular Phylogeny of *Orthochirus fomichevi* Kovařík, Yağmur, Fet & Hussen, 2 (0S1c9o rpiones: Buthidae) From Duhok Province, Kurdistan Region, Iraq, as Inferred from Mitochondrial *COI* Gene



Farhad Ramadhan Ali^{1*} , Hamid Saeid Kachel¹

1. Department of Biology, College of Science, University of Zakho, Zakho, Iraq.



How to cite this article Ali FR, Kachel HS. The First Molecular Phylogeny of *Orthochirus fomichevi* Kovařík, Yağmur, Fet & Hussen, 2 (0S1c9o rpiones: Buthidae) From Duhok Province, Kurdistan Region, Iraq, as Inferred from Mitochondrial *COI* Gene. *Archives of Razi Institute Journal*. 2025; 80(5):1179-1188. <https://doi.org/10.32598/ARI.80.5.3741>

doi <https://doi.org/10.32598/ARI.80.5.3741>

Article info:

Received: 12 Mar 2025

Accepted: 15 Jun 2025

Published: 01 Sep 2025

Keywords:

Buthidae, Genetic variation, Mitochondrial DNA, Phylogeny.

ABSTRACT

Introduction: *Orthochirus fomichevi* Kovařík, Yağmur, Fet & Hussen, 2019 comprises small-sized scorpions in the family Buthidae, inhabiting the Kurdistan region of Iraq and parts of Turkey. Despite the availability of some molecular data, the phylogenetic relationship and genetic variation within this species remain largely unexplored.

Materials & Methods: This study aimed to use the mitochondrial cytochrome c oxidase subunit I (*COI*) gene to build the phylogenetic relationships and estimate the evolutionary divergence of *O. fomichevi* collected from Duhok Province, Kurdistan Region, Iraq.

Results: A maximum likelihood tree of *O. fomichevi* revealed that this species forms a highly supported clade, distinct from other *Orthochirus* species. In contrast, *O. innesi* isolates from GenBank did not group under the same clade; instead, each formed a separate clade. Additionally, the *O. afghanus* isolate from GenBank failed to form a distinct clade and grouped with *O. persa* isolates from Iran. Mean genetic distances, calculated using the Kimura 2-parameter model, indicated that all analyzed sequences of *O. fomichevi* are genetically homogeneous and show greater similarity to *O. innesi* than to other species. Morphological analysis was in agreement with the molecular findings, further supporting the distinctiveness of *O. fomichevi*. In contrast, the *O. innesi* isolates were genetically highly divergent from each other, indicating that they belong to different taxa. Furthermore, *O. afghanus* and *O. persa* isolates from GenBank were genetically very close to each other, indicating that they belong to the same species.

* Corresponding Author:

Farhad Ramadhan Ali

Address: Department of Biology, College of Science, University of Zakho, Zakho, Iraq.

Tel: +964 (750) 1431025

E-mail: farhad.ali@staff.uoz.edu.krd



Copyright © 2025 The Author(s);
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).
Noncommercial uses of the work are permitted, provided the original work is properly cited.

Conclusion: This study presents the first molecular phylogeny of *O. fomichevi*, confirming its distinct genetic identity within the genus *Orthochirus* and highlighting potential cryptic diversity and misidentifications in *O. innesi*, *O. afghanus*, and *O. persa* isolates from the GenBank database. These findings underscore the importance of integrating molecular tools with morphological analysis for accurate scorpion taxonomy.

1. Introduction

O*rthochirus* Karsch, 1892 is a genus of scorpions within the family Buthidae. The genus was originally established by Karsch [1] as a replacement for the preoccupied name *Orthodactylus* Karsch, 1881. Since its inception, the taxonomic status of *Orthochirus* has been subject to considerable debate. Kraepelin [2] initially regarded it as a mere synonym of *Butheolus*, but it was later reinstated as a valid genus by Simon [3].

The genus *Orthochirus* comprises small-sized scorpions with a broad distribution across North Africa, the Middle East, Arabian Peninsula, Central Asia, and India [4]. To date, 53 species within this genus have been described [5], including three species recorded in Iraq: *O. fomichevi*, *O. mesopotamicus*, and *O. iraqus* [6]. According to Kachel et al. [7], each *Orthochirus* species in Iraq exhibits a restricted geographic range: *O. fomichevi* inhabits the northern region and the foothills of the Zagros Mountains; *O. iraqus* is distributed across the central and western plains; and *O. mesopotamicus* occurs in the southern humid and plain areas of the country. This spatial segregation underscores the species-specific habitat preferences and limited distributional overlap within Iraq.

O. fomichevi was first described as a new species based on morphological characteristics by Kovařík et al. [6]. Prior to this, it had been misidentified as *O. zagrosensis*; however, *O. fomichevi* can be differentiated by the presence of granules on the dorsal surface of metasomal segment V, a feature absent in *O. zagrosensis* [6]. The known distribution of *O. fomichevi* includes the Kurdistan region of Iraq and parts of Turkey. This initial misidentification underscores the limitations of relying exclusively on morphological traits for accurate species delimitation within the genus *Orthochirus*, highlighting the need for integrative taxonomic approaches.

DNA barcoding has emerged as a widely adopted molecular approach to overcome challenges in species identification, particularly for taxa exhibiting high morphological similarity or compromised specimen integrity. This technique primarily targets the 5' region of

the mitochondrial cytochrome c oxidase subunit I (*COI*) gene, which exhibits sufficient interspecific variability to serve as a standardized genetic marker for species-level discrimination, commonly referred to as the “barcode of life” [8, 9]. The effectiveness of DNA barcoding is further enhanced when integrated with detailed morphological analyses [10], providing a robust framework for accurate taxonomic resolution across diverse animal groups.

Although the mitochondrial *COI* gene sequence for *O. fomichevi* is available in the GenBank database, no published studies have yet addressed its phylogenetic placement within the genus. This gap highlights the necessity for comprehensive molecular analyses to elucidate the evolutionary relationships of *O. fomichevi*. Accordingly, this descriptive, cross-sectional study integrates morphological identification with mitochondrial *COI* gene analysis to infer the molecular phylogeny of *O. fomichevi*, contributing to a deeper understanding of its taxonomic and evolutionary context within *Orthochirus*.

2. Materials and Methods

2.1. Study area

The study area was conducted in Duhok Province in the Kurdistan region of Iraq, a predominantly mountainous region bordering Turkey and Syria (Figure 1). Geographically, it lies between latitudes 36°10'N and 37°23'N and longitudes 42°20'E and 44°19'E. Duhok Province is administratively divided into seven districts: Zakho, Sumel, Duhok, Amedi, Shekhan, Akre, and Bardarash. The overall study was conducted from June 2024 to March 2025. Ethical considerations were strictly adhered to throughout the study to ensure compliance with relevant guidelines and regulations.

2.2. Taxon sampling

O. fomichevi specimens were collected nocturnally between June and October 2024 using a high-intensity ultraviolet (UV) lamp, capitalizing on the species' nocturnal behavior and natural fluorescence under UV light, which facilitates detection. Collected specimens were divided for preservation based on subsequent analyses:

Some were fixed in 95% ethanol and stored at -20 °C for molecular studies, while others were preserved in 80% ethanol and maintained at room temperature for morphological examination. The geographical distribution of *O. fomichevi* was mapped using ArcGIS Pro software, version 3.0.1 (Esri, USA).

2.3. Species identification and imaging

All specimens were identified morphologically using the taxonomic key provided by Kovařík et al. [6]. Morphological identifications were verified by an experienced scorpion taxonomist. Morphological terminology follows the standards established by Sissom [11]. High-resolution images of whole specimens were captured using a Canon EOS 7D camera. Image stacking was performed using Helicon Focus software (HeliconSoft, Ukraine) to enhance depth of field. The focus stacking methodology was adapted from the Canon-Cognisys system recommended by Brecko et al. [12].

2.4. DNA extraction

Genomic DNA was extracted from approximately 20 mg of metasomal muscle tissue using the AddPrep Genomic DNA Extraction Kit (Addbio, Korea). Prior to extraction, muscle tissue was washed in physiological saline solution (PiONEER, Iraq) for 15–20 minutes to remove residual ethanol and subsequently dried on filter paper. The dried tissue was transferred to a 1.5 mL Eppendorf tube and homogenized using a propylene micropestle (Shenzhen Baimai Life Science, China). The subsequent extraction steps were conducted according to the manufacturer's protocol. The isolated DNA was stored at -20 °C until further analysis.

2.5. Mitochondrial COI gene fragment amplification

Polymerase chain reaction (PCR) amplification of the barcode region of the mitochondrial *COI* gene was performed using a TC1000-G-Pro Thermal Cycler (DLAB, China). The reaction employed the universal primers LCO1490 (5'-GGTCAACAAATCATCATAAAGATATTGG-3') and HCO2198 (5'-TAAACTTCAGGGTGACCAAAAAATCA-3') as described by Folmer et al. [13]. PCR reactions were conducted in a total volume of 20 µL, containing 0.5 µL (10 pmol) of each primer, 10 µL of Addstart Taq Master Mix (2× concentration; Addbio, Korea), 2 µL of template DNA (~20 ng/µL), and 7 µL of water. The thermal cycling conditions are detailed in Table 1. Amplification success was verified by electrophoresis on a 1.5% aga-

rose gel. PCR products were subsequently sent to Macrogen Inc. (Seoul, Korea) for Sanger sequencing.

2.6. DNA sequence processing

Sanger sequencing trace files were manually checked and trimmed by SeqTrace software, version 0.9.0 [14] to generate consensus sequences, which are then subjected to BLAST tool [15] against GenBank records to confirm DNA sequence identities and to search for closely related sequences for phylogenetic construction. The study sequences were deposited in GenBank under the accession numbers listed in Table 2.

2.7. Phylogenetic analysis

All sequences were aligned using the ClustalW algorithm implemented in MEGA software, 11 VERSION 11 [16]. Phylogenetic relationships were inferred using the maximum likelihood (ML) method, with branch support assessed via 1,000 bootstrap replicates. The optimal substitution model was selected based on bayesian information criterion (BIC) scores [17]. The resulting ML tree generated in MEGA 11 was exported in Newick format and subsequently visualized and edited using figtree software, version 1.4.4 to produce a high-quality phylogenetic tree.

Mean genetic distances within and between *O. fomichevi* and closely related sequences (excluding the out-group) were calculated using the Kimura 2-parameter model (K2P) [18] to estimate evolutionary divergence among the sequences.

3. Results

3.1. Taxonomy

3.1.1. Family Buthidae C. L. Koch, 1837

3.1.1.1. Genus *Orthochirus* Karsch, 1892

3.1.1.1.1. *O. fomichevi* Kovařík, Yağmur, Fet & Hussen, 2019

Type locality and type depository: Iraq, Sulaymaniyah Province, Chaqzhi Khwaroo; František Kovařík, private collection, Prague, Czech Republic (FKCP).

Materials examined: Iraq, Duhok province, Zakho district, Shahida, 37°08'01.1"N 42°40'24.0"E, 506 m above sea level (ASL), 2♂2♀; Khrababk, 37°08'30.6"N 42°45'02.1"E, 526 m ASL, 1♂3♀; Salka, 37°06'50.8"N 42°39'05.7"E, 573 m ASL, 2♂3♀; Betas, 37°04'02.6"N

Table 1. Steps and conditions of thermal cycling for PCR

Steps	Temperatures	Time	Cycles
Initial denaturation	95 °C	5 minutes	1 cycle
Denaturation	94 °C	30 seconds	35 cycles
Annealing	48 °C	30 seconds	
Extension	72 °C	30 seconds	
Final extension	72 °C	7 minutes	1 cycle




Figure 1. Geographical distribution of *O. fomichevi* plotted on the topography of Duhok Province

a) Map of Iraq showing the Kurdistan region (red and yellow polygons), with Duhok Province shown in red; b) Topographic map of Duhok Province: Blue and red circles indicate collection sites of the isolates used in morphological and molecular studies, respectively.

42°42'06.1"E, 675 m ASL, 1♂; Dubanik, 37°02'30.2"N 42°50'19.9"E, 709 m ASL., 4♂3♀; Sumel district, Sumel, 36°50'31.3"N 42°51'52.8"E, 446 m ASL, 6♂2♀; Esmahil Ava, 36°57'25.3"N 42°38'12.9"E, 504 m ASL, 13♂11♀; Duhok district, Nizarke, 36°50'20.7"N 43°04'03.5"E, 659 m ASL, 6♂7♀; Shekhan district, Atrush, 36°51'07.7"N 43°21'54.3"E, 1014 m ASL, 7♂8♀.

Table 2. Details of *O. fomichevi* isolates from Duhok Province used in molecular study, including their corresponding *COI* gene data

Locality Data				COI Gene Data	
Isolates	Localities	Latitude	Longitude	Accession Numbers	Length (bp)
DUH01	Esmahil Ava	36°57'25.3"N	42°38'12.9"E	PV219459	641
DUH02	Nizarke	36°50'20.7"N	43°04'03.5"E	PV219460	641
DUH03	Salka	37°06'50.8"N	42°39'05.7"E	PV219461	639
DUH04	Atrush	36°51'07.7"N	43°21'54.3"E	PV219462	639
DUH05	Atrush	36°51'07.7"N	43°21'54.3"E	PV219463	639




Figure 2. *O. fomichevi*, showing dorsal (a, c) and ventral (b, d) aspects of male (a, b) and female (c, d)

Detailed images: e) Sternite VII; f) Ventral aspect of metasomal segments I-III; g) Dorsal aspect of metasomal segment V and telson; scales 10 mm (a-c), 2 mm (e-g)

Abbreviations: mtI-V: Metasomal segments I-V; ti: Telson.

3.2. Diagnosis

The general body coloration is predominantly black, with the exception for the chelae fingers and the distal segments of the legs, which are yellow (Figure 2a, 2b, 2c and 2d). Adult body length ranges from 30.12 to 39.91 mm in males and 30.37 to 45.34 mm in females; with females generally larger than males. The trichobothrium d_2 on the femur is typically absent. The movable finger of the chela bears 8–9 rows of denticles, including both internal and external denticles, as well as four subtermi-

nal denticles. The number of pectinal teeth ranges from 21 to 23 in males and 17 to 20 in females. Mesosomal sternite VII presents a granulate surface characterized by prominent granulate carinae (Figure 2e). Metasomal segments II and III lack granules on the ventral and lateral surfaces, displaying a punctate and bumpy texture (Figure 2f). Metasomal segment V is distinguished by dense granulation along the dorsal mesial surface (Figure 2g). The telson is elongated and smooth, lacks granules. The vesicle is long, with lateral and ventral punctation. The aculeus is long and curved.

Table 3. Accession numbers of *Orthochirus* species and outgroup sequences retrieved from GenBank

Species	Localities	Accession Numbers
<i>O. fomichevi</i>	Iraq	MT232527
<i>O. innesi</i>	Morocco	ON255632
<i>O. innesi</i>	Egypt	MZ669861
<i>O. innesi</i>	-	JQ514244
<i>O. persa</i>	Iran	PP574852
<i>O. persa</i>	Iran	PP574854
<i>O. persa</i>	Iran	PP574853
<i>O. bicolor</i>	India	KT716038
<i>O. bicolor</i>	India	MF422331
<i>O. glabrifrons</i>	Oman	ON255631
<i>O. afghanus</i>	Afghanistan	ON255630
<i>O. vachoni</i>	Somaliland	ON255629
<i>H. tamulus</i>	Outgroup	KT716031

3.3. Phylogenetic analysis

The barcode region of the *COI* gene was successfully sequenced from five *O. fomichevi* specimens (Table 2), collected from four distinct localities. These sequences were supplemented with 12 additional *COI* sequences of various *Orthochirus* species retrieved from the GenBank database. Additionally, a sequence of *Hottentotta*

tamulus was included as an outgroup for phylogenetic analyses (Table 3).

A phylogenetic tree for *O. fomichevi* was constructed using the ML method, applying the TN93+G model [19], selected as the best-fit model by MEGA software. The ML tree with the highest log likelihood value of -1,738.62 was selected for visualization. In this tree (Fig-

Table 4. Estimates of net evolutionary divergence in the *COI* sequences within and between *Orthochirus* species

Row	Species	1	2	3	4	5	6	7	8	9
1	<i>O. fomichevi</i>	0.0054								
2	<i>O. innesi</i> (a)	0.0656	N/A							
3	<i>O. innesi</i> (b)	0.074	0.0745	N/A						
4	<i>O. innesi</i> (c)	0.1012	0.1091	0.0989	N/A					
5	<i>O. persa</i>	0.0933	0.0911	0.0863	0.1024	0.0067				
6	<i>O. glabrifrons</i>	0.1119	0.1315	0.1286	0.1451	0.1238	N/A			
7	<i>O. afghanus</i>	0.1122	0.1006	0.0902	0.1173	0.0237	0.1249	N/A		
8	<i>O. bicolor</i>	0.1159	0.1363	0.1084	0.1132	0.1131	0.118	0.1115	0.002	
9	<i>O. vachoni</i>	0.1268	0.1312	0.1325	0.1454	0.1457	0.1251	0.1491	0.1528	N/A

Note: The mean genetic distances calculated using the K2P model. The values in bold represent intraspecific divergence. (a), (b) and (c) represent the isolates mentioned in Figure 3.




Figure 3. Maximum likelihood phylogenetic tree of *O. fomichevi* inferred from the COI gene

Note: Study sequences are shown in red. Numbers above and below branches indicate bootstrap support values. The scale bar reflects the number of substitutions per nucleotide site. (a), (b), and (c) represent the isolates mentioned in Table 4.

ure 3), sequences of *O. fomichevi* formed a highly-supported clade with *O. fomichevi* from Iraq, as evidenced by a high bootstrap support value of 98. However, interspecific relationships among other *Orthochirus* species could not be well resolved due to low bootstrap values. In contrast, different *O. innesi* isolates did not form a clade with each other but instead each formed a separate clade, which indicates that they belong to different taxa. Additionally, *O. afghanus* did not form a separate clade but instead grouped with *O. persa* in a highly-supported clade (bootstrap value=98), indicating that they belong to the same taxon.

An examination of evolutionary divergence, calculated using the K2P model (Table 4), showed that the mean intraspecific divergence among members of *O. fomichevi* was low (0.54%), indicating a high level of genetic similarity within the species. Additionally, *O. fomichevi* exhibited a close genetic affinity to *O. innesi* from Morocco, with a mean interspecific divergence of 6.56%. In contrast, *O. fomichevi* was more genetically divergent from *O. vachoni*, with a divergence of 12.68%, compared to other *Orthochirus* species. Moreover, *O. innesi* isolates ON255632, JQ514244, and MZ669861 were highly divergent from each other, with divergence values

ranging from 7.45 to 10.91%. Additionally, *O. afghanus* was very close to *O. persa* (divergence value=2.37%).

4. Discussion

In the present study, eighty-one specimens of *O. fomichevi* were collected from multiple locations across Duhok Province. These scorpions were predominantly encountered in open plain habitats characterized by sparse vegetation. Morphological analysis consistently confirmed species identification based on diagnostic traits, while molecular analysis of the *COI* gene revealed that *O. fomichevi* forms a strongly supported clade (bootstrap=98) distinct from other *Orthochirus* species. The intraspecific divergence was low (0.54%), indicating high genetic homogeneity, while interspecific divergence with other species ranged from 6.56% to 12.68%. Notably, sequences labeled as *O. innesi*, *O. persa*, and *O. afghanus* in [GenBank](#) showed evidence of misidentification.

Previous records have documented *O. fomichevi* in the provinces of Duhok, Erbil, and Sulaymaniyah within the Kurdistan region of Iraq, as well as in Hakkari Province from Turkey [6, 20, 21]. Morphologically, *O. fomichevi* can be distinguished from other *Orthochirus* species by several diagnostic features, including a densely granulated mesosomal sternite VII with well-developed granulate carinae; a metasomal segment V exhibiting dense granulation along the dorsal mesial surface; and metasomal segments II and III that are smooth, punctate, and bumpy ventrally and laterally, lacking granules [6]. All examined specimens in this study exhibited these defining morphological characteristics.

To complement morphological identification, phylogenetic analysis based on the mitochondrial *COI* gene was performed. The resulting phylogenetic tree demonstrated that the specimens collected in this study and previously sequenced isolates of *O. fomichevi* form a strongly supported clade (bootstrap value=98), distinct from other *Orthochirus* species. Estimates of genetic divergence corroborate these findings; the mean intraspecific divergence within *O. fomichevi* was 0.54%, reflecting high genetic similarity among conspecific individuals. In contrast, mean interspecific divergence between *O. fomichevi* and other *Orthochirus* species ranged from 6.56% to 12.68%, with *O. innesi* from Morocco (ON255632) exhibiting the lowest divergence, thereby representing the closest relative to *O. fomichevi*.

These results are consistent with previous molecular studies on the genus *Orthochirus*, such as Jafari et al.

[22], who analyzed *COI* and *16S rRNA* gene sequences from *O. iranensis*, *O. farzanpayi*, *O. zagrosensis*, and *O. stockwelli* populations in Iran. Their findings similarly reported low intraspecific divergence alongside relatively higher interspecific divergence, supporting the phylogenetic patterns observed in the present study. However, they did not deposit their data in [GenBank](#); that's why we did not include them in our analysis. Similarly, Barahoei [23] also reported low intraspecific divergence, particularly in *O. persa*. The *Orthochirus* fauna of Iraq includes two other species: *O. mesopotamicus* and *O. iraqus* [7]. In this study, we were unable to infer their phylogenetic relationships due to the unavailability of their *COI* data in [GenBank](#).

Additionally, the three isolates of *O. innesi* (ON255632, JQ514244, and MZ669861) retrieved from [GenBank](#) did not group together; instead, each formed a separate clade. Genetic divergence analysis further supported these findings, in which the divergence between them was relatively high (7.45–10.91%). The phylogenetic relationships between the isolates JQ514244 (unknown) and MZ669861 (Egypt) were previously inferred by Mohammed-Geba et al. [24], in which the two isolates grouped under the same clade. However, their study was based on a wide range of divergent species from different genera, and no *Orthochirus* species other than the ones mentioned; therefore, they were grouped together. The type locality of *O. innesi* is Cairo, Egypt [3]; therefore, we conclude that the isolate MZ669861 represents the true *O. innesi*, while the isolates ON255632 and JQ514244 belong to two misidentified specimens which could not be assigned due to limited *COI* data in [GenBank](#).

Furthermore, *O. afghanus* from Afghanistan failed to form a distinct clade but instead formed a highly supported clade (bootstrap value=98) with the *O. persa* isolates from Iran. This indicates that they belong to the same species. The interspecific divergence further supports these findings, in which the divergence value between them was very low (2.37%), considered two separate species. The localities of the isolates of *O. afghanus* (ON255631) and *O. persa* (PP574852, PP574854, and PP574853) analyzed in this study are close to each other [23, 25]; therefore, it was not possible to conclude whether they belong to *O. afghanus* or *O. persa*. Generally, species misidentification is very common in the case of scorpions, even in dedicated literature [26], which has led to many incorrect placements of scorpions in the [GenBank](#) database [27].

A limitation of this study was the inability to comprehensively investigate certain districts due to the political instability, their distance from our research center, and the logistical challenges associated with sample collection in these areas. Additionally, the limited availability of *COI* sequence data for *Orthochirus* species in [GenBank](#) restricted comparative molecular analyses.

5. Conclusion

In conclusion, this study provides robust evidence for the accurate identification and phylogenetic placement of *O. fomichevi* within the genus *Orthochirus*, confirmed by consistent morphological traits and a strongly supported phylogenetic clade with low intraspecific genetic divergence. The unexpected divergence and placements of isolates labeled *O. innesi*, *O. persa*, and *O. afghanus* highlight the need for thorough taxonomic revision and expanded genetic sampling. To address challenges from misidentifications in public databases, we recommend integrating morphological and molecular tools in scorpion taxonomy and expanding molecular datasets to include additional *Orthochirus* species, which will improve database accuracy and enhance biodiversity knowledge in the region

Ethical Considerations

Compliance with ethical guidelines

The study protocol was approved by the Animal Ethics Committee of College of Science, [University of Zakho](#), Zakho Iraq (Code: AEC-031).

Data availability

The *COI* sequence data are available in [GenBank](#) at under the accession numbers PV219459, PV219460, PV219461, PV219462, and PV219463.

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' contributions

Study conceptualization and study design: All authors; Data acquisition, interpretation, statistical analysis and writing the original draft: Farhad Ramadhan Ali; Supervision, review and editing: Hamid Saeid Kachel.

Conflict of interest

The authors declared no conflicts of interest.

Acknowledgements

The authors would like to express their sincere gratitude to Ersen Aydın Yağmur ([Manisa Celal Bayar University](#), Turkey) for kindly photographing the specimens used in this study and for his valuable assistance in confirming their identification. Special appreciation is also extended to the Biology Research Center at the [University of Zakho](#) for kindly granting permission to conduct this research on their premises.

References

- [1] Karsch F. Arachniden von Ceylon und von Minikoy gesammelt von den Herren Doctoren P. und F. Sarasin. *Berl Entomol Z.* 1891; 36(2):267-310. [\[Link\]](#)
- [2] Kraepelin K. Pedipalpi. In: Dahl F, editor. *Das tierreich, 8 (Arachnoidea)*: Herausgegeben von der deutschen zoologischen gesellschaft. Berlin: R. Friedländer und Sohn Verlag; 1899. [\[Link\]](#)
- [3] Simon E. Révision des scorpions d'Égypte. *Bulletin de la société entomologique degypte.* 1910; 2:57-87. [\[Link\]](#)
- [4] Kovařík F, Just P. *Orthochirus katerinae* sp. n. (Scorpiones: Buthidae) from Saudi Arabia. *Euscorpius.* 2022; 2022(362):1-9. [\[Link\]](#)
- [5] Rein JO. *The Scorpion Files.* Trondheim: Norwegian University of Science and Technology; 2025. [\[Link\]](#)
- [6] Kovařík F, Yağmur EA, Fet V, Hussen FS. A review of *Orthochirus* from Turkey, Iraq, and Iran (Khoozestan, Ilam, and Lorestan Provinces), with descriptions of three new species (Scorpiones: Buthidae). *Euscorpius.* 2019; 2019(278):1-31. [\[Link\]](#)
- [7] Kachel HS, Al-Khazali AM, Hussen FS, Yağmur EA. Checklist and review of the scorpion fauna of Iraq (Arachnida: Scorpiones). *Arachnol Mitt.* 2021; 61(1):1-10. [\[DOI:10.30963/aramit6101\]](#)
- [8] Hebert PD, Ratnasingham S, deWaard JR. Barcoding animal life: Cytochrome c oxidase subunit 1 divergences among closely related species. *Proc Biol Sci.* 2003; 270(Suppl_1):S96-9. [\[DOI:10.1098/rsbl.2003.0025\]](#) [\[PMID\]](#)
- [9] Sheir SK, Galal-Khallaf A, Mohamed AH, Mohammed-Geba K. Morphological and molecular clues for recording the first appearance of *Artemia franciscana* () in Egypt. *Heliyon.* 2018; 4(12):e01110. [\[DOI:10.1016/j.heliyon.2018.e01110\]](#) [\[PMID\]](#)

- [10] Mohammed-Geba K, Sheir SK, El-Aziz Hamed EA, Galal-Khallaf A. Molecular and morphological signatures for extreme environmental adaptability of the invasive mussel *Brachidontes pharaonis* (Fischer, 1870). *Mol Cell Probes*. 2020; 53:101594. [DOI:10.1016/j.mcp.2020.101594] [PMID]
- [11] Sissom WD. Systematics, biogeography, and paleontology. In: Polis GA, editor. *The biology of scorpions*. Stanford: Stanford University Press; 1990. [Link]
- [12] Brecko J, Mathys A, Dekoninck W, Leponce M, Vanden-Spiegel D, Semal P. Focus stacking: Comparing commercial top-end set-ups with a semi-automatic low budget approach. A possible solution for mass digitization of type specimens. *Zookeys*. 2014; (464):1-23. [DOI:10.3897/zookeys.464.8615] [PMID]
- [13] Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol*. 1994; 3(5):294-9. [PMID]
- [14] Stucky BJ. SeqTrace: A graphical tool for rapidly processing DNA sequencing chromatograms. *J Biomol Tech*. 2012; 23(3):90-3. [DOI:10.17171/jbt.12-2303-004] [PMID]
- [15] Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990; 215(3):403-10. [PMID]
- [16] Tamura K, Stecher G, Kumar S. MEGA11: Molecular evolutionary genetics analysis version 11. *Mol Biol Evol*. 2021; 38(7):3022-7. [DOI:10.1093/molbev/msab120] [PMID]
- [17] Nei M, Kumar S. *Molecular evolution and phylogenetics*. Oxford: Oxford University Press; 2000. [DOI:10.1093/oso/9780195135848.001.0001]
- [18] Kimura M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol*. 1980; 16(2):111-20. [DOI:10.1007/BF01731581] [PMID]
- [19] Tamura K, Nei M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol*. 1993; 10(3):512-26. [PMID]
- [20] Hussen F, Kachel H, Hama G, Kachal E, Slo M, Hiwil I, et al. Epidemiological characterizations, new localities, and a checklist of the known scorpions in the Kurdistan Region, Northern Iraq. *J Arthropod Borne Dis*. 2022; 16(3):251-61. [DOI:10.18502/jad.v16i3.12042] [PMID]
- [21] Hussen FS, Ahmed ST. New data of scorpion fauna, include two new records with identification key of scorpion species (Arachnida: Scorpiones) in Iraq. *Plant Arch*. 2020; 20(2):6711-25. [Link]
- [22] Jafari H, Salabi F, Forouzan A. A study of genetic diversity among different population of *Orthochirus* sp. based on cytochrome C oxidase subunit I and 16srRNA sequencing. *Turk J Zool*. 2020; 44(1):64-8. [DOI:10.3906/zoo-1904-9] [PMID]
- [23] Barahoei H. Unlocking the genetic diversity of Scorpions (Arachnida: Scorpiones) in a lowland area of Sistan: A DNA barcoding study. *Taxon Biosyst*. 2024;16(61):9-14. [DOI:10.22108/tbj.2025.143533.1289]
- [24] Mohammed-Geba K, Obuid-Allah AH, El-Shimy NA, Mahbob MA, Ali RS, Said SM. DNA Barcoding for Scorpion Species from New Valley Governorate in Egypt Reveals Different Degrees of Cryptic Speciation and Species Misnaming. *Conservation*. 2021; 1(3):228-40. [DOI:10.3390/conservation1030018]
- [25] Štundlová J, Šťáhlavský F, Opatová V, Stundl J, Kovařík F, Dolejš P, et al. Molecular data do not support the traditional morphology-based groupings in the scorpion family Buthidae (Arachnida: Scorpiones). *Mol Phylogenet Evol*. 2022; 173:107511. [DOI:10.1016/j.ympev.2022.107511] [PMID]
- [26] Lourenço WR. Scorpion incidents, misidentification cases and possible implications for the final interpretation of results. *J Venom Anim Toxins Incl Trop Dis*. 2016; 22:1. [DOI:10.1186/s40409-016-0075-6] [PMID]
- [27] Soltan-Alinejad P, Rafinejad J, Dabiri F, Onorati P, Terenius O, Chavshin AR. Molecular analysis of the mitochondrial markers COI, 12S rDNA and 16S rDNA for six species of Iranian scorpions. *BMC Res Notes*. 2021; 14(1):40. [DOI:10.1186/s13104-021-05449-3] [PMID]